

STRAITS BRANCH  
ROYAL ASIATIC SOCIETY

---

---

[No. 41]

**JOURNAL**

January, 1904

Agents of the Society

London: KEGAN PAUL, TRENCH, TRUBNER & Co.

u

nd

4

[No. 41]

JOURNAL  
of the  
Straits Branch  
of the  
Royal Asiatic Society

JANUARY 1904

---

SINGAPORE:  
PRINTED AT THE AMERICAN MISSION PRESS  
1904





## Table of Contents.

---

Council for 1904 ... ..	v
Proceedings of Annual General Meeting ... ..	vi
Annual Report of the Council ... ..	viii
Treasurer's Cash Account for 1903 ... ..	ix
Obituary ... ..	x

---

Two Sea Dyak Legends, by the <i>Rev. Edwin H. Gomes</i> ...	1
New Malayan Plants, by <i>H. N. Ridley</i> ... ..	31
Notes on a Cruise in the Southern China Sea, by <i>C. Boden</i> <i>Kloss</i> ... ..	53
A List of the Butterflies of Borneo, by <i>R. Shelford</i> ...	81
The Sakais of Batang Padang, Perak, by <i>G. B. Cerruti</i> ...	113
On some Hymenoptera from the Raffles Museum, Singa- pore, by <i>P. Cameron</i> ... ..	119
Short Notes ... ..	125



THE  
STRAITS BRANCH  
OF THE  
ROYAL ASIATIC SOCIETY.

---

*COUNCIL FOR 1904.*

The Right Rev. BISHOP HOSE, *President.*

Hon. C. W. S. KYNNEERSLEY, C.M.G., *Vice-President  
for Singapore.*

Hon. DR. W. C. BROWN, *Vice-President for Penang.*

H. N. RIDLEY, Esq., M.A., *Honorary Secretary.*

DR. R. HANITSCH, *Honorary Treasurer.*

P. J. BURGETS, Esq., M.A.

Hon. W. R. COLLYER, M.A., I.S.O.

H. ESCHKE, Esq.,

REV. W. G. SHELLABEAR,

W. G. ST. CLAIR, Esq.,

}  
} *Councillors.*  
}

## Annual Report for 1903.

The Council have the pleasure to state that the financial position of the Society continues satisfactory, as may be seen by the Treasurer's Balance Sheet.

The number of members at present is 145, including the following gentlemen elected during the year.

DR. ABBOTT,	MR. F. C. MARSHALL,
MR. ERIC MAXWELL,	" R. D. HUDSON,
" GEORGE MAXWELL,	" F. W. DOUGLAS,
" W. H. CRADDOCK,	" W. S. GIBSON,
" A. H. BURN MURDOCH,	" T. C. HINKS,
" E. W. BIRCH,	" HON. H. F. DESHON,
" W. MAKEPEACE	" REV. H. C. IZARD,
" A. S. HAYNES,	" S. MOORHOUSE,

They have to express their great regret at the loss by death of Mr. A. W. O'Sullivan, long a member of the Society and at one time the Secretary, and also of Mr. D. H. Wise and Mr. James Driver.

Only one Journal No. 39 was published during the year but another will be shortly in the hands of the members. An important article by Mr. R. Shelford of Sarawak Museum on the Musical Instruments of Borneo illustrated by a large number of plates will follow as soon as it can be printed.

Referring to the resolution passed at the general meeting last year to procure and publish Malay Manuscripts, the Council desire to state that they have kept the matter in view, but up to the present no Manuscripts considered worthy of publication have as yet been obtained. Two important Manuscripts however have recently been offered to the Society by Mr. George Maxwell, and it is hoped they may appear in the course of this year.

The Library was rearranged and is being catalogued. Many books, journals and pamphlets were received and a number were bound.

During the year a letter was received from the Royal Asiatic Society of Bengal stating that the members of that Society had decided that the members of the Straits Branch of the Royal Asiatic Society should have the right of admission to the Society's meetings whenever they were in Calcutta.

The Society has now attained its twenty-fifth year having been founded on Nov. 4, 1877, and it may be noted that in spite of early prophecies of its soon becoming extinct it has steadily thrived to the present day.

The Treasurer's statement of accounts is appended.

HONORARY TREASURER'S ACCOUNT FOR THE YEAR 1903.

	\$	c.	\$	c.
Balances brought forward from 1902:—				
Mercantile Bank, Fixed Deposit	2395	54		
Chartered Bank, Fixed Deposit	700	81		
Mercantile Bank, Current Account	348	10		
Chartered Bank, Current Account	589	5		
			3703	40
Receipts in 1903:—				
Subscriptions for 1901	10	...		
do. " 1902	50	...		
do. " 1903	510	...		
do. " 1904	25	...		
do. for Life Memberships	100	...		
Sale of Journals	310	25		
Sale of Maps	962	21		
Sundry Recoveries	350			
Bank Interest	1207	1	2091	67
			5795	07
Payments in 1903:—				
American Mission Press	414	50		
Thacker and Spink	273	7		
Mounting of Maps and Binding of Books	315	25		
3 New Book-cases, at \$85 each	255	...		
Clerk's Salary	93	15		
Peon's Salary	15	...		
Postages and Petties	78	24		
			1198	51
Balances carried forward:—				
Mercantile Bank, Fixed Deposit	2700	...		
Chartered Bank, Fixed Deposit	1000	...		
Mercantile Bank, Current Account	549	29		
Chartered Bank, Current Account	347	27	4596	56
			5795	07

Examined and found correct.

A. KNIGHT.

R. HANITSCH.

*Honorary Treasurer, Straits Branch, Royal Asiatic Society.*

## OBITUARY.

### Mr. Arthur W. S. O'Sullivan.

Since the date of the last report the Society has had to lament the loss of Mr. Arthur W. S. O'Sullivan, a member of our Council for several years, and at one time Secretary.

Mr. O'Sullivan was born in 1860 and after a distinguished career at Trinity College, Dublin (scholar and gold medallist), he entered the Straits Settlements Civil Service in 1883.

Throughout his service of twenty years in this Colony he was distinguished as an able hard-working officer and showed a marked talent for languages—he was proficient in Dutch, Tamil, Malay, and more than one dialect of Chinese, which is a record rare amongst Europeans in this climate. He had held the post of Assistant Colonial Secretary for five years and had just been selected by the Colonial Office for the post of Colonial Secretary at Trinidad when he was struck down after a brief illness. Although he was not a frequent contributor to the *Journal*, the Society has lost in him an intellectual force—a man who took a keen interest in scholarship of every kind. For three years before his death he was engaged in intervals of leisure in the translation of Dr. Snouck Hurgronje's *Acheen*—a valuable and interesting piece of work which will shortly be published at Leiden in Holland.

In endeavoring to open up the wide field of Dutch learning and experience in Netherlands India to English readers, he has set an example for which the Society may well be grateful.

*R. N. Bland.*



## Two Sea-Dyak Legends.

By the REV. EDWIN H. GOMES, M.A.

There are many fairy tales and legends known to the Sea-Dyaks of the present day. These seem to be handed down, by word of mouth, from generation to generation from ancient times.

These stories may be roughly divided into two classes:—

I. Those which are purely fabulous and related as such, and are simply meant to interest and amuse, and in these respects resemble the fairy tales familiar to us all; and

II. Those which are believed to be perfectly true, and to have actually taken place, and are the traditions respecting their gods and preternatural beings. These form in fact the Mythology of the Dyaks.

To the first class belong a large collection of stories corresponding to the Adventures of Brer Fox and Brer Rabbit. In the Dyak tales, the *Plandok* and the *Kekura* (the mouse deer and the tortoise) act always in concert, and their combined intelligence is victorious over the rest of the animal world. To this class also belong the numerous stories related of *Apai Samumang* the Dyak type of cunning and wiliness—and *Apai Saloi*—the typical Dyak fool.

To the second class belong the many and varied adventures of *Klieng*, the great hero of ancient times, and his wife *Kumang*, the Dyak Venus, as well as the traditions relating to the gods believed in by the Dyaks of the present day. To these must be added certain stories which give a reason for some of the curious customs observed by the Dyaks. The two Dyak Myths which follow belong to this latter class.

## I.

**Danjai and the Were-Tiger's Sister.**

Once upon a time there lived a great Chief named *Danjai*. He was the head of one of the longest Dyak houses that were ever built. It was situated on a hill in the midst of a large plantation of fruit trees. *Danjai* was said to be very rich indeed. He possessed much farming land, many fruit trees, many *Tapang* trees, where the wild bees make their abode, and from which the sweet honey is obtained, and in his room there were many valuable jars of various kinds, and also a large number of brass vessels; for the Dyaks convert their wealth into jars and brassware to hand down to posterity. Every year he obtained a plentiful harvest of paddy much more than he and his family could consume and he had always much paddy for sale, so much so that the news of his wealth travelled to distant lands, and many from afar off would come and buy paddy from him. *Danjai* also possessed many slaves who were ready to help him in his work. All the people in his house had a very high opinion of his judgment, and were ready to obey his decisions, whenever he settled any of their disputes. So great indeed was his reputation for wisdom, that men from distant villages would often consult him and ask his advice when in any difficulty. He had also great fame as a brave warrior, and during expeditions against the enemy, he was the leader of the men of his own village and of many villages around, for all liked to follow such a brave man as *Danjai*, who was sure to lead them to victory. Over the fireplace in his verandah he had, hanging together in a bunch, the dried heads of the enemies whom he himself had killed.

Now this man *Danjai* had a very pretty wife whom he had recently married, but the marriage feast had not been held, because he had not yet obtained a human head from the enemy as a token of his love for her: for this girl was of a good birth and a Chief's daughter and wanted the whole world to learn, when they attended her marriage feast, what a brave man her husband was.

*Danjai* said to his young wife, "I will hold a meeting of the Chiefs around, and tell them that we must all get our war-boats ready, as I intend leading an expedition against the enemy.



I should like to bring you a human head as a token of my love, so that you may not be ashamed of your husband. And as soon as I return, we will have the wedding feast."

And though his wife was sorry that her husband intended leaving her, still she did not oppose his wishes, for she wished him to come back covered with glory.

So a council of war was held, and *Danjai* told the assembled Chiefs what he intended to do, and it was decided that all should begin at once making war-boats, which were to be ready in two months' time.

*Danjai* assisted by his slaves and followers, had been at work at his boat for several weeks, and it was nearly finished. It was a beautiful boat made out of the trunk of one large tree, and *Danjai* was proud of his work. He was so anxious to finish his boat, that one day he started very early in the morning, before his breakfast was ready, and he asked his wife to bring his food to him later on to the part of the jungle where he was working at his boat.

So Mrs. *Danjai* cooked the food and then ate her own breakfast. Then she made up small bundles of rice and also put together some fish and salt, and placed all in a little basket to take to her husband. She had never been out in the jungle by herself before, but she was not afraid, for her husband had told her the way, and she could hear the sound of his adze as he worked at his boat not very far off. She hung her basket over her left shoulder and, holding her small knife in her right hand, went cheerfully on. Presently she came to the stump of a tree on which was placed a bunch of ripe rambutan fruit. They looked so tempting that she could not help eating some of them, and as they were very nice, she put what remained in her basket, saying to herself, "Perhaps *Danjai* forgot to take these fruits with him and left them here. I will take them to him myself, he will no doubt be glad to eat these ripe fruits after his hard work."

Now there was in that land a Were-Tiger, that was much feared by all who lived around. He had the appearance of a man, but at times would transform himself into a tiger, and then he would attack human beings and carry off their heads as trophies to his own house. But he never attacked any unless

they had first done wrong by taking something which belonged to him. So this Were-Tiger would leave tempting fruit by the side of jungle paths, and on the stumps of trees, in the hope that some tired traveller would take and eat them. And if any one ate such fruit, then he or she was doomed to be killed by him that same day. But all knew about him, and though he placed many tempting baits in all parts of the jungle, no one touched his fruit, for all feared the fate which awaited them if they did any such thing. But *Danjai's* wife knew nothing about the Were-Tiger. No one had told her of him, and she had never been out before in the jungle by herself, and she had never been warned not to touch any fruit she might find lying about.

"Oh *Danjai*," she said, as soon as she met her husband, "I am afraid I am rather late. You must be very tired and hungry, working the whole morning at your boat without having had anything to eat. Never mind! Here is your breakfast at last." And she handed the basket which contained his food to her husband.

Now *Danjai* was really very hungry, so he was glad to see his food had arrived. He thanked his wife and at once began to empty the basket.

The first thing he saw was the ripe rambutan fruit at the top, and he asked his wife where she got them from. She told him she had found them on the stump of a tree by the wayside, and she said she thought they had been left there by him. She added with a smile, that they were very good as she had eaten some herself.

Then *Danjai*, brave man though he was, turned pale with fear and anxiety.

"We must not linger here a moment," he said to his wife. "Hungry though I am, I will not eat my food here. We must both hurry home at once. You have taken and eaten fruit belonging to the Were-Tiger, so much feared by all. It is said that whoever touches his fruit will surely die a terrible death: and you are the first person I know who has done so."

*Danjai* hurriedly gathered together all his tools and told those that were with him of his trouble, and they all started and walked silently back. *Danjai* was wondering how he was to



avert the fate which awaited his young wife. She was silent, because she saw her husband was troubled, and she was sorry that she had caused him grief.

As soon as they arrived at the house, *Danjai* sent for all the men round about and told them what had happened, how his wife had taken and eaten the fruit of the Were-Tiger. He begged them all to help to shield her, for the Were-Tiger was sure to have his revenge, and come and take the head of his wife.

So they all prepared themselves for the tiger's visit by sharpening their knives and spears. Some men placed themselves on the roof of the house, others in the verandah. The ladder leading up to the house was also guarded, and so were all parts of the house by which he was likely to force an entrance. As for *Danjai's* wife, they hid her beneath some mats and sheets in the room, and twelve brave men stood round her with their swords drawn, ready to save her life even at the cost of their own.

Just before dark they heard the roar of the tiger in the distance. Though still a long way off, the sound was very terrible to hear, and the men all grasped their swords and spears firmly, for they knew the tiger would soon be upon them.

Once more the tiger's roar sounded, nearer and clearer, and then they heard him crash through the leaf thatch roof and fall into the room. There was a great commotion among the men, but though all tried to kill the animal, none could see him. Soon after they heard a roar of triumph from the tiger outside the house. They lifted up the mats and sheets which covered *Danjai's* wife, and there they saw her headless body! The Were-Tiger had succeeded in his attack, and had carried off the head of his victim!

Loud was the weeping and great the lamentation over her dead body. She was so young to die! And what death could be more terrible than hers whose head had been carried away by her murderer! All in the house mourned her loss for seven days and during that time the house was very quiet, as all lived in their separate rooms, and did not come out into the common verandah to do work or to talk to each other.

The death of his wife grieved *Danjai* very much. But though his grief was great, his desire for revenge was greater still.

Very early on the morning of the next day, *Danjai* started after the tiger. The drops of blood which had fallen could plainly be seen on the ground, and he had no difficulty in finding out in what direction the tiger had gone. On and on he tracked the blood till he came to a cave at the foot of a high mountain. The sides of the cave were splashed with blood, so *Danjai* walked boldly in, determined to revenge the death of his wife. It was not very dark in the cave. In the distance he could see an opening and he hurried towards it.

He came out on the other side of the mountain, and saw a large plantation of sugar-cane and plantain trees. Beyond this he saw a long Dyak house.

"This," he said to himself, "is surely the abode of the Were-Tiger, and soon I shall have an opportunity of revenging the death of my wife."

He planted two sticks across one another in the ground to mark the opening in the mountain, so that he might not miss his way on his return, and then he boldly walked towards the house.

He followed a path through the sugar-cane plantation—still tracking the drops of blood upon the ground—until he came to the ladder leading up to the house. He was so anxious to attack his wife's murderer, that he did not pause to ask—as is the usual Dyak custom—whether he might walk up or not, but went straight on into the house. Men sitting in the verandah asked him, as he passed them, where he was going and what he wanted, but he did not answer them. His heart was heavy within him, thinking of his dead wife, and wondering whether he would be able to accomplish his task, and whether he would succeed in leaving the house as easily as he came in. But he was determined to avenge his wife's murder, and he would not shrink from any difficulties in the way.

He stopped at the room of the head of the house, and a girl asked him to sit down, and spread a mat for him. He did so, and the girl went into the room to fetch the brass vessel containing the betelnut ingredients which the Dyaks love to chew. As he sat down, he saw drops of blood on the fire-place, and looking up he noticed a fresh head, still dripping with blood, among the other skulls hanging there. He recognised it at a glance—it was the head of his loved wife!!



The girl came out with the brass vessel of betelnut and said: "Help yourself *Danjai*. We did not expect you to visit us so soon. Please excuse me for a little while, I have to attend to the cooking. But you will not be alone for my brother will soon be back. He has only gone to the plantation to fetch some sugar-cane."

So *Danjai* sat on the mat by himself, thinking what he was to do next and what he was to say to his wife's murderer when he came in. Soon the Were-Tiger arrived, carrying on his shoulder a bundle of sugar-cane.

"I am very pleased to see you *Danjai*," he said, "would you like some sugar-cane? If so, help yourself."

*Danjai* was so sad thinking of his wife, that he did not notice how curious it was, that they should know his name when they had never seen him before. He did not feel at all inclined to eat sugar-cane, but lest his host should think he had come to kill, and to put him off his guard, he pretended to eat a little. He heard the Were-Tiger say to his sister in the room, that she was to be sure to have enough food cooked, as *Danjai* would eat with them that evening. Then he left them and went to the river to bathe.

The sister came out of the room, and spoke to *Danjai*, who was still sitting in the verandah, and asked him to come into the room as she had something to say to him.

"Yes, *Danjai*," she said to him in a kind tone of voice, "I know of your trouble and I am sorry for you. However, if you follow my advice, all will be well. You must be careful, for my brother is easily put out, and has no scruples about killing any who displease him. Even our own people here hate him, for he is so merciless; but no one dare attack him, for all fear him greatly. Now listen attentively to what I have to say. When I put out the plates of rice in the room presently, do not take the one he tells you to have: take any of the others, for the one he wishes you to have is sure to contain some poison. Later on, when you retire to rest, do not spend the night on the mat spread out for you, but sleep somewhere else, and put the wooden mortar for pounding paddy on the mat in your stead: and so again on the second night, place the wooden mill for husking the paddy on your mat: and on the third night a roll of

the coarse matting used for treading paddy. If his three attempts to kill you are unsuccessful, then he will be in your power and will do what you command. But even then there is still danger, and you must not do anything rash, but ask my advice again later on. But go outside now into the verandah, for I think I hear my brother returning from his bath. I must make haste and put out the food for you all to eat."

Soon the Were-Tiger came in and sitting on the mat by *Danjai* asked him the news and how matters were in his country. *Danjai* answered little for he was very sad, besides his host always laughed at him whenever he spoke. The fact was that he was amused at the idea of the man, whose wife he had killed, sitting in his verandah and talking to him in a friendly way.

The sister came out of the room and asked them in to have their meal. All happened as she said it would. *Danjai* remembered her advice and did not take the plate of rice his host offered him. But he was too sad to eat.

In the evening *Danjai* and the Were-Tiger sat by a fire in the verandah. Over this fire hung several human heads. The tears came into *Danjai's* eyes as he sat there and saw the head of his dear wife being scorched by the fire. He felt inclined there and then to grasp his sword and attack the murderer of his wife; but he restrained himself remembering the advice of the Tiger's sister.

The Were-Tiger said to him with a nasty laugh, "What is troubling you that you should weep?"

"I am not troubled about anything," said *Danjai*, "but the smoke of the fire is too much for my eyes, and it makes them water and feel sore."

"If so," said his host, "let us put out the fire and retire to rest, as it is very late."

Two mats were spread out for them, one on each side of the fire-place, and they lay down to sleep. But *Danjai* kept awake, and when his companion was asleep, he rose and placed the wooden mortar for pounding paddy on his mat, and covered it over with a sheet; and he himself retired to a safe place as he was advised to do by the Tiger's sister. He watched to see what would happen and he was not disappointed. Not long after, he saw the Were-Tiger wake up and fetch a sword, and walk



up to the place where he was supposed to be asleep. With the sword he made two or three vicious cuts at the wooden mortar and said:

"Now *Danjai*, this will settle you. You will not think of revenging yourself on me any more."

Then *Danjai* cried out from where he was, "What is the matter? What are you doing?"

"Oh *Danjai*! Is that you?" said his host, "I did not mean to hurt you. I had a bad dream, and I sometimes walk in my sleep. How lucky it is you were not lying on the mat! I should have certainly killed you, and I should never have forgiven myself for doing so. Please understand I meant no harm to you, and let us lie down to rest again."

On the two following nights the Were-Tiger attempted to kill *Danjai*, but failed each time, because following the advice given him, *Danjai* placed first the wooden mill for husking the paddy on his mat, and next a roll of coarse matting used for treading paddy. His host made the same excuse for his strange behaviour each time.

On the morning of the fourth day, after the Were-Tiger had left the house to see whether any fish had been caught in his fish trap, his sister asked *Danjai* to come into the room as she had something to say to him before he left to return home.

"Now *Danjai*," she said, "as I told you before, since my brother has not been able to kill you these three days, he is in your power. After breakfast ask him to accompany you and show you the way back to your country. When you have both come to the further end of the sugar-cane plantation, ask him to sit down for a little while, and say you would like to eat some sugar-cane, before you leave him and go on your journey alone. When he gives you the sugar-cane, ask him to lend you his sword, giving as an excuse that yours is not sharp enough for peeling the sugar-cane, or that it is stuck fast in its sheath and cannot be drawn. When he hands you his sword, you must attack him with it and kill him. My brother is invulnerable to any other sword but his own. When you have killed him, cut off his head and bring it to me, and I will give you your wife's head in exchange for it. On no account are you to take his head away with you. If you do so, I will follow you to your country and take my revenge."

A few minutes after this conversation, the Were-Tiger returned with a basket full of fish. Some of these were soon cooked, and they sat down to breakfast.

Soon after they had eaten, *Danjai* told his host that he must be returning to his own country, and asked him to accompany him and show him his way back. So they started together and walked through the sugar-cane plantation.

Just as they came near the end of it, *Danjai* asked his companion to stop. He said he would like to have some sugar-cane before going on.

"I am sorry I did not offer you any," said the Were-Tiger: "it was very forgetful of me. Never mind, I will at once cut down some sugar-cane for us."

When he had brought the sugar-cane and had finished peeling the piece he wanted for himself, *Danjai* said to him,

"Please lend me your sword, for mine is stuck fast in its sheath and I cannot draw it out."

The Were-Tiger suspecting nothing, handed the sword to him, and *Danjai* began peeling his sugar-cane.

Just then the Were-Tiger turned round to look at his house, and *Danjai* seizing his opportunity, gave him a blow with the sword in his hand and killed him. Then he cut off the head and carried it back with him to the house he had just left.

When he came near, he saw the sister watching for his return, and standing at the top of the ladder leading up to the house. He followed her into the house, and gave her the head of her brother.

"You ought to be quite satisfied now, *Danjai*," she said, "for you have killed my brother, and have taken your revenge for the death of your wife. I want you to promise me certain things before you go. First of all, you must not let anybody know that you have killed my brother. Next, on your return, you must go on the war-path and bring back to me the head of a woman, to enable me to put away the mourning of myself and my relatives, for the death of my brother. And when you return, I hope you will take me with you to be your comforter in the place of your dead wife: so that I may have some one to care for me, now that my brother is dead. And I give you now some locks of my hair, to be used as a charm to



make you invisible to the enemy, when you are on the warpath. Lastly, I advise you and your people, never to eat or to take away any fruit you may find lying about in the jungle, on the stump of a tree, or on a rock, without knowing for certain who put it there and to whom it belongs, or making sure that it has fallen from some tree near. This must be remembered from generation to generation. Whoever disobeys this advice will be punished either by death, as in the case of your wife, or in some other dreadful way. You may now have the head of your wife to take back to your country; and as you may have forgotten the way, I will send one of my slaves with you, to show you in what direction you are to go."

As she finished speaking, she handed him his wife's head, and *Danjai* started off at once for he was anxious to get back.

He reached his house late that same evening. All his friends were glad to see him come back safe and sound. They had given up all hope of seeing him again. They were also pleased to see he had been successful in bringing back the head of his dead wife.

Soon after *Danjai's* return from the Were-Tiger's country he gathered all his followers together and told them that he intended going on the war-path. As soon as they were able to get everything ready, they started for the enemy's country. They were very successful and succeeded in taking many heads; but *Danjai*, protected as he was by the charm which he had received from the Were-Tiger's sister, was more successful than the others. They returned with much rejoicing, and a great feast was held in honor of their victory. The human heads were placed on a costly dish, and the women carried them into the house, with dancing and singing.

A few days after, *Danjai* started to fulfil his promise to the Were-Tiger's sister. He brought her back with him as his wife, and they lived very happily together for many years.

This story explains why the Dyaks, even at the present day, dare not eat any fruit they may find lying on the stump of a tree, or on a rock in the jungle. They fear that evil will happen to them, as it did to *Danjai's* wife.

## II.

## The Story of Siu,

Who first taught the Dyaks to plant Paddy and to observe the Omens of Birds.

Many thousands of years ago before the Paddy plant was known, the Dyaks lived on tapioca, yams, potatoes and such fruit as they could procure. It was not till *Siu* taught them how to plant Paddy that such a thing as rice was known. The story of how he came to learn of the existence of this important article of food, and how he and his son *Seragunting* introduced it among their people is here set forth.

*Siu* was the son of a great Dyak chief, but his father died when he was quite a child, and at the time this story begins, he lived with his mother and was the head of a long Dyak house in which lived some three hundred families. He was strong and active and handsome in appearance, and there was no one in the country round who was equal to him in strength or comeliness. When he was ready to go on the warpath, he was the admiration of all the Dyak damsels. On such occasions he appeared in a many coloured waistcloth, twelve fathoms in length, which was wound round and round his body. On his head was a plaited rattan band in which were stuck some long feathers of the hornbill. His coat was woven of threads of bright colours. On each well-shaped arm was an armlet of ivory. To his belt was fastened his sword and the many charms and amulets that he possessed. With his spear in his right hand and his shield on his left arm, he presented a splendid type of a Dyak Warrior. But not of his bravery nor of his deeds of valour against the enemy does this tale relate. It only gives an account of an adventure of his which ended in his discovery of Paddy.

One day *Siu* proposed to the young men of his house that they should take their blowpipes with them and go into the jungle to shoot birds. So one morning they all started early. Each man had with him his bundle of food for the day, and each went a different way, as they wished to see, on returning in the evening, who would be the most successful of them all.

*Siu* went towards a mountain not far from his house. He wandered about the whole morning in the jungle, but strange to



say, he did not see any bird nor did he meet with any animal. Everything was very quiet and still. Worn out with fatigue, he sat down to rest under a large tree, and feeling hungry, he ate some of the food he had brought with him. It was now long past midday, and he had not been able to kill a single bird! Surely none of the others could be so unfortunate as he!

He determined not to be beaten by the others, and after a short rest, he started again and wandered on in quest of birds. The sun had gone half-way down in the western heaven, and he was beginning to lose heart, when suddenly he heard not far off the sound of birds. He hurried in that direction, and came to a large wild fig tree covered with ripe fruit, which a large number of birds were busy eating. Never before had he seen such a sight! On this one large tree, the whole feathered population of the forest seemed to have assembled together! On looking carefully, he was surprised to see that the different kinds of birds were not all intermingled together as is usually the case. Each species was apart from the others. Here he saw a large flock of wild pigeons on one branch, and next to them were the parrots, all feeding together but keeping distinct from them. Upon this tree there were hornbills, wood-peckers, wild pigeons and all the different kinds of birds he had ever seen.

He hid himself under the thick leaves of a shrub growing near, very much pleased at his luck. He took a poisoned dart and placed it in his blow-pipe, and taking good aim, shot it out. He had aimed at one bird in a particular flock, and he hit it. But that bird was not the only one that fell dead at his feet. To his astonishment, he saw that many of the other birds that were near it were killed also. Again he shot out a dart, and again the same thing happened. The bird that was hit fell down dead, and with it the birds that were near it. In a very short time, Siu had killed as many birds as he could carry. As the little basket, in which he had brought his food, was too small to hold them all, he set to work and made a large coarse basket with the bark of a *Pendok* tree growing near. Then he put his load on his back and started to return home, glad that he had been so successful.

He tried to follow the way by which he had come, but as he had not taken the precaution to cut marks in the trees he passed,

he very soon found himself in difficulties. He wandered about, sometimes passing by some large tree, which he seemed to remember seeing in the morning. He climbed up a steep hill and went several miles through a large forest, but did not find the jungle path which he had followed early in the day. It was beginning to grow dusk and the sun had nearly set.

"I must hurry on," said *Siu* to himself, "in the hope of finding some house where I can get food and shelter. Once it is dark, I shall be forced to spend the night in the jungle."

He hurried on and luckily came to a part of the jungle which had lately been a garden.

"There must be some path from this garden," said *Siu* to himself, "leading to some house;" and he began to walk round it.

He soon found an old disused path which he followed and which led him to another path. By this time it was quite dark, and *Siu* made haste to reach the Dyak house which he felt sure was not very far off. He soon came to a well, and not far off he saw the lights and heard the usual sounds of a Dyak house. He was glad to think that he would not have to spend the night in the jungle, but would be probably able to get food and shelter at the house.

He stopped to have a bath and hid the birds he was carrying and his blow-pipe and quiver in the brushwood near the well, hoping to take them with him when he started to return the next morning.

As he approached the house, he could hear the voices of the people there. When he came to the bottom of the ladder leading up to the house, he shouted, "Oh! you people in the house, will you allow a stranger to walk up?" At once there was dead silence in the house. No one answered. Again *Siu* asked the same question, and, after a pause, a voice answered, "yes: come up!"

He walked up into the house. To his surprise, he saw no one in the open verandah in front of the different rooms. That part of a Dyak house, usually so crowded, was quite empty. Nor did *Siu* hear the voices of people talking in any of the rooms. All was silent. Even the person who answered him was not there to receive him.



He saw a dim light in the verandah, further on, in the middle of the house, and he walked towards it. He wondered what could have happened to all the people in the house, for not long before he heard many voices.

"This seems to be a strange house," he said to himself. "When I was bathing and when I walked up to the house, it seemed to be well inhabited, but now that I come in, I see no one, and hear no voice."

When *Siu* reached the light, he sat down on a mat there. Presently he heard a woman's voice in the room say, "Sit down *Siu*: I will bring out the *pinang* and *sirih* to you."

*Siu* was very pleased to hear a human voice. Soon a young and remarkably beautiful girl came out of the room with the chewing ingredients, which she placed before him.

"Here you are at last, *Siu*," she said, "I expected you would come earlier. How is it you are so late?"

"I stopped a little while at the well to have a bath, as I was hot and tired."

"You must be very hungry as well," she said, "wait a moment while I prepare some food for you. After you have eaten we can have our talk together."

When *Siu* was left to himself, he wondered what it all meant. Here was a long Dyak house, built for more than a hundred families to live in, and yet it seemed quite deserted. The only person in it appeared to be the beautiful girl who was cooking his food for him. Then again, he wondered how it was she knew his name and expected him that day. All these things filled him with wonder and surprise.

"Come in, *Siu*," said the voice from the room, "your food is ready."

*Siu* was very hungry and went in at once, and sat down to eat his dinner.

When they had done eating, she cleared away the plates and put things back into their places and tidied the room. Then she spread out a new mat for him, and brought out the *pinang* and *sirih*, and bade him be seated, as she wished to have a chat with him.

*Siu* had many questions to ask, and as soon as they were both seated, he began:—

"Why are you all alone in this house? This is a long house, and many families must live in it; where are the others? Why is everything so silent now? I am sure I heard voices before I entered the house; but now I hear no sound."

"Do not let us talk about this house or the people in it for the present. I would much rather talk of other matters. Tell me of your own people, and what news you bring from your country."

"There is no news to give you," *Siu* replied. "We have been rather badly off for food, as our potatoes and yams did not turn out so well this year as we hoped."

"Tell me, what made you come in this direction and how it was you found out this house."

"While I was hunting in the jungle to-day, I lost my way. After wandering about a long time, I found a path which I followed and came to this house. It was kind of you to take me in and give me food. If I had not found this house, I must have died in the jungle. To-morrow morning I must ask you to show me the way to my country, and also I must beg of you some food for my journey back. My mother is sure to be anxious about me. She is left all alone, now that I am away. My father died a long time ago, and I am her only son."

"Do not go away as soon as to-morrow morning. Stay here a few days at any rate."

At first *Siu* would not consent, but she spoke so nicely to him that she succeeded in persuading him to stay there at least a week. Then he went out to the verandah, and she brought out a mat for him to sleep on and a sheet to cover himself with. As *Siu* was very tired, he soon fell sound asleep, and did not wake up till late on the following morning.

He saw some little children playing about the next day, but he did not see any grown up people. He went into the room to have his morning meal, but saw no one there, except the girl he had seen the evening before. He felt very much inclined to ask her again where the people of the house were, but he did not do so, as she did not seem inclined to speak about them.

Now though *Siu* knew it not, this was the house of the great *Singalang Burong*, the Ruler of the Spirit World. He was able to metamorphose himself and his followers into any



form. When going forth on an expedition against the enemy, he would transform himself and his followers into birds, so that they might travel more quickly. Over the high trees of the jungle, over the broad rivers, sometimes even across the sea *Singalang Burong* and his flock would fly. There was no trouble about food, for in the forests there were always some wild trees in fruit, and while assuming the form of birds, they lived on the food of birds. In his own house and among his own people *Singalang Burong* appeared as a man. He had eight daughters, and the girl who was cooking food for *Siu* was the youngest of them.

The reason why the people of the house were so quiet, and did not make their appearance, was because they were all in mourning for many of their relatives who had been killed some time back. Only the women and children were at home, because that same morning all the men had gone forth to make a raid upon some neighbouring tribe, so that they might bring home some human heads to enable them to end their mourning. For it was the custom that the people of a house continued to be in mourning for dead relatives, until one or more human heads were brought to the house. Then a feast was held, and all mourning was at an end.

After *Siu* had been in the house seven days, he thought he ought to think of returning to his own people. By this time he was very much in love with the girl who had been so kind to him, and he wished above all things to marry her, and take her back with him to his own country.

"I have been here a whole week," he said to her, "and though you have not told me your name, still I seem to know you very well. I have a request to make and I hope you will not be angry at what I say."

"Speak on; I promise I will not be angry whatever you may say."

"I have learnt to love you very much," said *Siu*, "and I would like to marry you if you will consent, so that I shall not leave you but take you with me, when I return to my own land. Also I wish you to tell me your name, and why this house is so silent, and where all the people belonging to it are."

"I will consent to marry you, for I also love you. But you must first promise me certain things. In the first place, you must not tell your people of this house and what you have seen here. Then also you must promise faithfully never to hurt a bird or even to hold one in your hands. If ever you break this promise, then we cease to be man and wife. And of course, you must never kill a bird, because if you do so, I shall not only leave you but revenge myself on you. Do you promise these things?"

"Yes" said *Siu*, "I promise not to speak of what I have seen here until you give me leave to do so. And as you do not wish it, I will never touch or handle a bird, and certainly never kill one."

"Now that you have promised what I wish, I will tell you about myself and the people of this house," said the maiden. "My name is *Endu-Sudan-Galinggam-Tinchin-Mas* (the girl Sudan painted like a gold ring), but my people call me by my pet names *Bunsu Burong* (the youngest of the bird family), and *Bunsu Katupong* (the youngest of the *Katupong* family). This house, as you noticed, seems very empty. The reason is that a month ago many of our people were killed by some of the people of your house, and we are all still in mourning for them. As you know, when our relatives have lately died, we stay silent in our rooms, and do not come out to receive visitors or to entertain them. Why are your people so cruel to us? They often kill our men when they go out fishing or hunting. On the morning of the day on which you arrived, all the men of this house went on the war-path, so as to obtain the heads of some of the enemy to enable us to put away our mourning. With us as with you it is necessary that one or more human heads be brought into the house, before the inmates can give up sorrowing for their dead relatives and friends. You see us now in the form of human beings, but all the people in this house are able to transform themselves into birds. My father *Singalang Burong* is the head of this house. I am the youngest of eight sisters: we have no brother alive. Our only brother died not long ago, and we are still in mourning for him, and that was the reason why my sisters did not come out to greet you."

*Siu* heard with surprise all she had to say. He said to himself that it was lucky he did not bring up to the house the



basket of birds which he had killed in the jungle, and that he had hidden them with his blow-pipe and quiver containing poisoned darts, in the brushwood near the well. He determined to say nothing about the matter, as probably some of her friends or relations were among the birds that were killed by him.

So *Siu* married *Bungu Burong* and continued to live in the house for several weeks.

One day he said to his wife:—"I have been here a long time. My people must surely be wondering where I am, and whether I am still alive. My mother too must be very anxious about me. I should like to return to my people, and I want you to accompany me. My mother and my friends are sure to welcome you as my wife."

"Oh yes: I will gladly accompany you back to your home. But you must remember and say nothing of the things you have seen in this house. When shall we start?"

"We can start early to-morrow morning, soon after breakfast," answered *Siu*.

They started early the next day, taking with them food enough for four days, as they expected the journey would last as long as that. *Siu's* wife seemed to know the way, and after journeying for three days, they came to the stream near the house, and they stopped to have a bath. Some of the children of the house saw them there, and ran up to the house and said:—" *Siu* has come back, and with him is a beautiful woman, who seems to be his wife."

Some of the older people checked the children, saying:—"It cannot be *Siu*: he has been dead for a long time. Don't mention his name, for if his mother hears you talk of him, it will make her very unhappy."

But the children persisted in saying that it was indeed *Siu* that they had seen. Just then *Siu* and his wife appeared, and walked up to the house.

*Siu* said to his wife:—"The door before which I hang up my sword is the door of my room. Walk straight in. You will find my mother there, and she will be sure to be glad to welcome you as her daughter-in-law."

When they came into the house, all the inmates rushed out to meet them, and to congratulate *Siu* on his safe return.

They asked him many questions:—where had he been living all this time; how he came to be married, and what was the name of his wife's country. But *Siu* answered little, as he remembered the promise he had made to his wife, that he would not speak of what he had seen in her house.

When they reached the door of his room, *Siu* hung up his sword and his wife went into the room. But she did not see his mother as she was ill and was lying in her curtain. Then *Siu* followed his wife into the room and called out "Mother, where are you? Her is your son *Siu* come back!"

But his mother made no answer, so he opened her curtain, and saw her lying down, covered up with a blanket. She had been so troubled at the thought that her son was dead, that she had refused to eat and had become quite ill.

She would not believe that her son had really returned alive, and she said, "Do not try to deceive me; my son *Siu* is dead."

"I am indeed your son *Siu*, and I have come back alive and well!"

"No!" she replied, "my son *Siu* is dead. Leave me alone, I have not long to live. Let me die in peace and follow my son to the grave."

*Siu* then went to the box in which his clothes were kept, and put on the things that his mother had often seen him wear. Then he went to her again and said, "Even if you do not believe that I am your son, at any rate you might turn round and look at me, to make sure that I am not your son."

Then she looked at him, and saw that it was indeed her son. She was so pleased at his return that she soon recovered from her illness, which was really caused by her sorrow and refusal to eat. *Siu* told his mother of his marriage, and she welcomed his wife with joy.

The women all crowded round *Siu's* wife and asked her what her name was. She answered *Endu-Sudan-Galingam-Tinchin-Mus*. (The girl Sudan painted like a gold ring). They looked at her in surprise; they had never heard of such a name before.

"Where do you come from?" they asked. "What is the name of your country?"



"*Nanga Niga Bekurong Bebuli ugadi Tekugong Mabong,*" (The mouth of the hidden Niga stream changed into the Mabong snail),\* was the reply.

They were astonished at her answer! They had never heard of such a country. They asked her of her people, but she would not say anything more of herself or speak about her people.

Everybody admired the great beauty of *Siu's* wife. No more questions were asked of her, as she seemed unwilling to answer. Her parentage remained a mystery.

In process of time *Siu's* wife bore him a son whom they named *Seragunting*. He was a fine child, and as befitted the grandson of *Singalang Burong*, he grew big and strong in a miraculously short time, and when he was three years old, he was taller and stronger than others four times his age.

One day as *Seragunting* was playing with the other boys, a man brought up some birds which he had caught in a trap. As he walked through the house, he passed *Siu* who was sitting in the open verandah. *Siu* forgetting the promise he had made to his wife asked him to show him the birds, and he took one in his hands and stroked it. His wife was sitting not far off, and saw him hold the bird and was very much vexed that he had broken his promise to her.

She got up and returned to her room. *Siu* came in and noticed that she was troubled and asked her what was wrong. She said that she was only tired.

She said to herself:—"My husband has broken his word to me. He was done the thing he promised me he would never do. I told him he was never to hold a bird in his hands, and that if he did such a thing, I would leave him. I cannot stay here in this house any longer. I must return to the house of my father *Singalang Burong*."

She took the water vessels in her hands, and went out as if to fetch water. But when she came to the well, she placed the water vessels on the ground, and disappeared in the jungle.

In the meantime *Seragunting*, tired with his play came back in search of his mother. She was very fond indeed of him, and

\*The Dyaks are fond of rhyming names, which often have no special meaning.

he expected her to come to him as soon as he called out to her. But he was disappointed. No one answered his call, and when he looked in the room, she was not there. He asked his father where his mother was, and he told him that she had just gone to the well to fetch water and would soon be back.

But hour after hour passed, and she did not return to the house. So *Seragunting* began to be anxious, and asked his father to accompany him to the well to look for her. At first his father refused to do so, but when he saw his son crying for his mother, he went with him to the well. They found the water vessels there, but saw no signs of her.

"Your mother is not here, *Seragunting*," said *Siu*. "Perhaps she has gone to the garden to get some vegetables for our dinner. Let us go back to the house. If your mother is not back, early to-morrow morning, we will go and look for her." So they both returned to the house, taking back with them the water gourds which *Siu's* wife had left at the well.

Early the next morning, *Seragunting* and his father went in search of her. They took with them only a little food, as they expected to find her not very far off. But they wandered the whole day and saw no signs of her. They spent the night under a large tree in the jungle. Early the next morning, they were surprised to find a small bundle of food, wrapped up in leaves, near *Seragunting*. This food was evidently meant for him alone, as it was not enough for two, but he gave some of it to his father, who ate sparingly of it, so that his son might not be hungry. They wandered on for several days, and every night the same strange thing occurred—a bundle of food was left near *Seragunting*. *Siu* suggested to his son that they should return; but *Seragunting*, who during the journey had grown up into a strong lad with a will of his own, would not consent to do so, as he was determined to find his mother.

They wandered on for several days, deeper and deeper into the jungle; but could find no signs of her whom they sought. At last they came to the sea-shore. Here they rested for some days, in the hope that some boat might pass. Still, as before, each morning a bundle of food was found by *Seragunting*. If it were not for this food, they would have long ago died of starvation. On this food they managed to live, waiting



hopefully to see some boat appear to take them on their journey.

One day as *Seragunting* was watching, he heard the sound of paddles, and saw in the distance several long boats approaching. He hailed the first, and asked the men in it to take him and his father with them. The boat made for the shore, but the man in the bows recognised the two wanderers, and shouted out:—"It is *Siu* and his son *Seragunting*: do not let them come into the boat." The boat went on and left them to their fate. The same thing happened in the case of each of the other boats. As soon as *Siu* and his son were recognised, no one would help them.

Now these were the boats of the sons-in-law of *Singalang Burong*:—*Katupong*, *Beragai*, *Bejampong*, *Papau*, *Nendak*, *Pangkak*, and *Embuas*. They were not pleased at their sister-in-law marrying a mere mortal like *Siu*, and so refused to help him and his son.

The next day *Seragunting* saw what seemed to be a dark cloud come towards him over the sea. As it came nearer, it took the form of a gigantic spider, carrying some food and clothes.

"Do not be afraid," said the Spider, "I have come to help you and your father. I have brought you food and clothing. When you have had some food and changed your clothes, I will take you across the water to the land on the other side. My name is *Emplawa Jawa* (the Spider of Java). I know your history, and I will lead you to your mother whom you seek."

After they had eaten and put on the new clothes brought them, the spider told them to go with him across the sea. They were not to be afraid, but to follow his track, not turning to the right hand nor to the left. They obeyed his words. Strange to say, the water became as hard as a sandbank under their feet. For a long time they were out of sight of land, but towards evening they approached the opposite shore, and saw a landing place where there were a large number of boats. Not far off where several houses, and one longer and more imposing than any of the others. To this house the Spider directed *Seragunting*, telling him that he would find his mother there. The Spider then left them. As it was late, they did not go up to

the house that evening, but spent the night in one of the boats at the landing place. Among the boats were those belonging to the Sons-in-law of *Singalang Burong* which had passed *Siu* and his son as they waited on the sea-shore for some boat to take them across the sea.

When *Seragunting* and his father woke up next morning, they saw that the road leading up to the house had sharpened pieces of bamboo planted close together in the path, to prevent their walking up it. As they were wondering what they were to do next, a fly came to *Seragunting* and said:—

“Do not be afraid to walk up. Tread on the spikes that I alight on; they will not hurt you. When you come to the house you will find swords with blades turned upwards fastened to the ladder. Tread on the blades that I alight on and walk boldly up into the house.”

They did as the fly advised them, and were not hurt. The bamboo spikes crumbled under their feet, and sword blades they trod on were blunt and harmless.

The people of the house took no notice of them, and they sat down in the verandah of the house. Then the fly came to *Seragunting* and whispered to him:—“You must now follow me into the room. Your mother is there, lying in her curtain. I will point out to you which it is, and you must wake her up and tell her who you are. She will be very pleased to see you. Then when you come out into the verandah and see the sons-in-law of *Singalang Burong*, you must greet them as your uncles. They will disown you and pretend that you are no relation of theirs. But do not be afraid. You will be victorious in the end.”

*Seragunting* followed the fly into the room and went to the curtain on which it alighted. He called out to his mother, and she awoke and saw with joy her son. She embraced him, and he said to her:—

“How is it you went away and left us? We missed you so much, and were so sorry to lose you, that my father and I have been travelling for many days and nights in search of you. Now our troubles are over for I have found you.”

“My dear son,” she said as she carressed him, “though I left you I did not forget you. It was I who placed the food by you



every night. I left your father because he broke the promise he made to me. But you are my own son, and I have been wishing to see you, ever since I left your house. It was I who sent the Spider to help you and show you your way here. My love for you is as great as it ever was. We will go out now into the verandah, and I will introduce you to your uncles and aunts and to your grandfather. They may not welcome you, because they were opposed to my marriage to your father. But do not be afraid of them. We will be more than a match for them all."

Then she spoke to her husband *Siu*, whom she was glad to meet again. All three then went out into the verandah, which was now full of people. *Seragunting* called the sons-in-law of *Singalang Burong* his uncles, but they refused to acknowledge that he was their nephew.

They proposed several ordeals to prove the truth of his words, that he was indeed the grandson of *Singalang Burong*. In all of these *Seragunting* came off victorious.

As the men and boys were spinning their tops, they asked *Seragunting* to join them. He had no top of his own, so he asked his mother for one. She took an egg and uttered some mysterious words over it, and immediately it became a top. This she gave to her son, who went and joined the others in the game. Whenever *Seragunting* aimed at a top, he always hit it and smashed it in pieces. None of the others were a match for him. In a short time, all the tops except that of *Seragunting* were broken in pieces.

Then they suggested a wrestling match. *Seragunting* was quite ready to try a fall with any of them, old or young. Some of their best wrestlers came forward. The first two were overthrown so easily by him, that the others saw it was no use their attempting to wrestle with *Seragunting*.

As a last trial they proposed that all should go out hunting. Here they hoped to be more fortunate. All the sons-in-law of *Singalang Burong* took their good hunting dogs with them, confident of success. *Seragunting* was told that he could have any of the other dogs left in the house. There he saw a few old dogs, weak and useless for hunting. With these he was expected to compete against the others, and if he was not successful, both he and his father were to be killed! *Seragunting*

consented even to such an unfair ordeal as that. He called to him an old sickly looking dog, and gently stroked it. At once it became young and strong! While the others went forth into the jungle with a pack of hounds, *Seragunting* was only accompanied by one dog. In the evening *Katupong*, *Beragai*, *Bejampong* and the others all returned unsuccessful. Soon after, *Seragunting's* dog appeared chasing a huge boar which made a stand at the foot of the ladder of the house. *Seragunting* asked the others to kill the beast if they dared. The spears cast at it glided off, and left the beast unharmed. Some of those who were rash enough to go near the animal, had a close escape from being torn in pieces by its tusks.

*Seragunting* armed with nothing better than a little knife belonging to his mother, walked up to the infuriated animal, and stabbed it in a vital part, and it fell down dead at his feet.

After these marvellous feats, all were compelled to admit that *Seragunting* was a true grandson of the great *Singalang Burong*. They all acknowledged him as such, and he was taken to his grand-father, who was pleased to see the lad and promised to help him throughout his life.

But *Siu* was unhappy in his new home. He could not help thinking of his mother whom he had left alone, and he was anxious to return to his own people. He begged his wife to accompany him back to his old home, but she refused to do so. It was decided that *Siu* and his son should stay in the house of *Singalang Burong* till they had obtained such knowledge as would be useful to them in the future, and that then they were to return to the lower world, bringing with them the secrets they had learnt from those wiser and more powerful than themselves.

All the people of the house were now most kind to *Siu* and his son, and were most anxious to teach them all they could. They were taken on a war expedition against the enemy, so they might learn the science and art of Dyak warfare. They were taught how to set traps to catch deer and wild pig. They were shown the different methods of catching fish, and learnt to make the different kinds of fish-trap used by the Dyaks of the present day. They remained in *Singalang Burong's* house that whole year, so that they might have a complete



and practical knowledge of the different stages of paddy growing.

When the year was ended, *Seraunting's* mother took him and *Siu* to see her father, *Singalang Burong*, so that they might receive from him his advice, as well as such charms as he might wish to give them before they left to return to the lower world of mortals.

*Singalang Burong* was sitting in his chair of state, and received them most kindly when they came to him. He bade them be seated on the mat at his feet, as he had many things to say to them. Then he explained to *Siu* and his son who he was, and the worship due to him, and they learnt also about the observance of omens, both good and bad.

"I am the Ruler of the Spirit World," said *Singalang Burong*, "and have the power to make men successful in all they undertake. At all times, if you wish for my help, you must call upon me and make offerings to me. Especially must this be done before you go to fight against the enemy, for I am the God of War and help those who pay me due respect."

"You have learnt here how to plant paddy. I will give you some paddy to take away with you, and when you get back to your own country, you can teach men how to cultivate it. You will find rice a much more strengthening article of food than the yams and potatoes you used to live upon, and you will become a strong and hardy race.

"And to help you in your daily work, my sons-in-law will always tell you whether that you do is right or wrong. In every work that you undertake, you must pay heed to the voices of the sacred birds:—*Katupong*, *Beragai*, *Bejompong*, *Papau*, *Nendak*, *Pangkas*, and *Embuas*. These birds, named after my sons-in-law, represent them and are the means by which I make known my wishes to mankind. When you hear them, remember it is myself speaking through my sons-in-law for encouragement or for warning. Whatever work you may be engaged in—farm work, house-building, fishing or hunting—wherever you may be you must always do as these birds direct. Whenever you have a feast, you must make an offering to me, and you must call upon my sons-in-law to come and partake of the feast. If you do not do these things, some evil is sure to happen to you. I

am willing to help you and to give you prosperity, but I expect due respect to be paid to me, and will not allow my commands to be disobeyed."

Then *Singalang Burong* presented them with many charms to take away with them. These were of various kinds. Some had the power to make the owner brave and fortunate in war. Others were to preserve him in good health, or to make him successful in his paddy planting and cause him to have good harvests.

*Siu* and *Seragunting* then bade their friends farewell, and started to return. As soon as they had descended the ladder of the house of *Singalang Burong*, they were swiftly transported through the air by some mysterious power, and in a moment they found themselves at the bathing place of their own house.

Their friends crowded round them, glad to see them back, safe and well. They were taken with much rejoicing to the house. Friends and neighbours were told of their return, and a great meeting was held that evening. All gathered round the two adventurers, who told them of their strange experiences in the far country of the Spirit Birds. The charms received from *Singalang Burong* were handed round for general admiration. The new seed paddy was produced, and the good qualities of Rice as an article of food explained. The people congregated there had never seen paddy before, but all determined to be guided by *Siu* and *Seragunting*, and to plant it in future. The different names of the Sacred Birds were told to the assembled people, and all were warned to pay due respect to their cries.

And so, according to the ancient legend, ended the old primitive life of the Dyak, when he lived upon such poor food as the fruits of the jungle, and any yams and potatoes he happened to plant near his house; the old blind existence, in which there was nothing to guide him; and then began his new life, in which he advanced forward a step, and learnt to have regularly, year by year, his seed-time and harvest, and to know that there were unseen powers ruling the Universe, whose will might be learnt by mankind, and obedience to whom would bring success and happiness.

## Note.

On receiving the above legends from Mr. Gomes I pointed out to him the curious fact that in the first legend the tiger—a beast quite unknown in Borneo—plays a prominent part, and suggested that this story may have been of Malay origin. He replies by referring to A. H. Everett's paper on the Tiger in Borneo, in Journal 5, p. 157, and says "the 'Tree-tiger' *Felis marmorata* is common enough. The Dyaks call it by a distinctive name 'Kemaung dau' or 'Kemaung raras' (dau and raras both being words meaning the branch of a tree). These would lead one to suppose that at one time they knew of some other species they called simply 'Kemaung.'" Everett refers to traditions of the animal also, among the Sea Dyaks. One may compare these traditions of an animal apparently absent from the country with those of the Mias (Mawas) of the Malay peninsula.

H. N. Ridley.





## New Malayan Plants.

By H. N. RIDLEY.

The following plants apparently undescribed have turned up lately in various collections made in the Peninsula.

### SCITAMINEÆ.

*Zingiber Wrayii*, Prain mss. A slender plant over a foot tall. Leaves elliptic, lanceolate, glabrous, 9 inches long, 3 inches wide, narrowed at the base but not petioled. Peduncle 4 inches tall covered with large loose sheathing leaves, spike three inches long, ovoid obconic. Bracts thin elliptic  $1\frac{1}{2}$  inch long  $\frac{1}{2}$  inch wide or less. Flowers solitary yellow. Bracteole spathaceous, lip spotted and marked with purple. Anther narrow linear, beak half an inch long.

Upper Perak at 300 feet elevation (Wray 3735). The only specimen I have seen is in poor condition but it seems a distinct plant from any described, from its rounded head of thin bracts, most of the allied species having cylindric spikes.

*Elettariopsis cyantescens*, n. sp. Rhizome  $\frac{1}{4}$  inch through, violet inside, covered with dry sheaths, stems 18 inches tall, rather slender base, olivaceous. Leaves 8, lanceolate acuminate glabrous 6 inches long  $1\frac{1}{2}$  inches wide, hardly petiolate, ligule rounded short. Spike short, peduncle half an inch long covered with long dry lanceolate bracts  $1\frac{1}{2}$  inches long, flowers four. Ovary oblong pubescent  $\frac{1}{4}$  inch long. Calyx tube  $1\frac{1}{2}$  inches long, pubescent at the base, apex long acuminate. Corolla tube narrow  $\frac{1}{2}$  inch longer pubescent, lobes oblong half an inch long  $\frac{1}{4}$  inch wide lip, oblong rounded 1 inch long. All white except the tip of the lip which is blue. Stamen oblong, crest moderately large, stigma large cup-shaped with pubescent edges.

This pretty species was collected by Mr. W. G. Napier on banks at Bukit Tanga, Seremban; it is nearly allied to *E. pubescens* but has much shorter flower spikes and quite glabrous leaves. The flowers when bruised became of a blue color, perhaps containing indigo.

#### ORCHIDEÆ.

*Dendrobium mellitum*, n. sp. A very small epiphytic plant, stems few 4 or 5 inches long, very slender above the lowest two or three joints, swollen a little for  $\frac{3}{4}$  inch length. Leaves few terete subulate, 3 inches long  $\frac{1}{16}$  inch thick. Flower solitary subterminal, from a short raceme, with small bracts. Pedicel and ovary  $\frac{1}{8}$  inch long thick yellow. Upper sepal ovate  $\frac{1}{4}$  inch long, lateral sepals ovate falcate obtuse, mentum short broad and blunt. Petals narrower and shorter, all yellow. Lip three-lobed, lobes broad short curved tips rounded yellow veined with pink inside, midlobe oblong truncate, edged minutely laciniate veins and a patch in the centre pink, three parallel raised veins white on the disc. Anther oblong hemispheric brown, column face flat stained pink.

Johore, at Castlewood on the Sungei Tebrau, May 1903, I found a single plant of this very weak slender little orchid on a tree, and flowered it in the Botanic Gardens. It is allied to *D. clavator*, Ridley, a native of Perak, but has the base of the stem much less swollen, and an entirely different lip. Considering the size of the flower, half an inch across, it was wonderfully strongly scented of honey.

*Thrixspermum crassifolium*, n. sp. An epiphyte on coffee bushes, stem 6 inches tall, thick, leaves close set, thick leathery oblong obtuse, tip rounded, 4 inches long  $1\frac{1}{2}$  inches thick, sheaths under half an inch long purple, scapes 3 to 5 inches long stiff, base purple, raceme  $1\frac{1}{2}$  to 4 inches long flattened. Bracts distichous  $\frac{1}{4}$  inch long close set, ovary and pedicel  $\frac{1}{4}$  inch long. Sepals linear caudate from a broader base  $\frac{3}{4}$  inch long, petals narrower linear caudate, all bright pellow. Lip  $\frac{1}{4}$  inch long, pubescent orange

passing into cherry red at the tip. Spur saccate blunt, side lobes arched blunt, middle fleshy blunt ovate. Callus on the disc tongue-shaped rounded depressed in the centre, column white with rounded and broad wings. Anther hemispheric. Fruit linear angled, 3 inches long  $\frac{1}{4}$  inch through.

On old coffee bushes at Castlewood, Johore. This is allied to *Th. Arachnites* which was common in the same place, but has the short stout leaves and stem of *Th. Scortechinii*. The flowers however are much smaller than any other of this group.

## AMARYLLIDÆ.

*Curculigo megacarpa*, n. sp. A large tufted plant with a stout stem. Leaves dark green oblong lanceolate acuminate glabrous 20 inches long, 3 inches wide, petiole 8 inches long. Peduncle 4 inches long tomentose with large lanceolate acuminate bracts at the base, lower ones hairy on the edge only, upper ones more hairy. Spike broad conic-cylindric 2 to 4 inches long. Flower 1 inch across. Sepals lanceolate acute  $\frac{1}{2}$  inch long dirty yellow outside and hairy at the tip. Petals bright yellow, stamens yellow, filaments short, anthers oblong, wavy, hairy, beaked. Fruit 2 inches long, Indian-club shaped  $\frac{3}{8}$  inch through at the base, white seeds very numerous angled ribbed black.

Perak, in forests on the Thaiping Hills at 2000 to 3000 feet elevation.

This seems to be common on the Thaiping Hills and has probably been hitherto overlooked or taken for a form of *C. latifolia*, but it can hardly be referred to that species, variable as it is. It is larger in all parts of the flower and fruit, the latter being very long club shaped.

## BURMANNIACEÆ.

*Burmannia oblonga*, n. sp. A slender saprophyte 5 inches tall with numerous wiry roots. Stem sparingly or not at all branched. Leaves sheath-like, scales few and distant. Flowers one or two on the ends of the stem  $\frac{3}{8}$  inch long and  $\frac{1}{4}$  inch broad across the wings, tube narrow wings oblong with straight edges, angles rounded. Sepals and petals short blunt. Petals much the shorter.



Penang on rocks at 1500 to 2000 feet altitude (King's collector 2270.)

This plant should be sought again. It resembles the common *B. coelestis* in the form of its flowers, but is saprophytic. The very broad oblong wings of the perianth tube are very striking.

Since publishing the paper on *Burmanniaceæ* in Journal 22, p. 332, I have been able to add to this group of plants:—*B. Championii*, Thwaites. A small ivory white plant like *B. tuberosa*, Becc., but more compact, with a short thick root stock covered with scales, from the Laba river in Selangor, and two additional species of *Thismia*; *Th. grandiflora*, Ridl., with rather large pink flowers collected by Lieut. Kelsall on the Sembrong river in Johor; and *Th. chrysops*, Ridl., a very pretty kind from Mount Ophir with pink and chocolate flowers with a yellow ring round the mouth of the tube; and *Bagnisia crocea* of Beccari, a very oddly shaped little brown species met with in the Perak Hills.

#### DIOSCOREACEÆ.

*Dioscorea tennifolia*, n. sp. Stems slender leaves mostly opposite thin glabrous elliptic cuspidate, base rounded 5 nerved 2 to 3 inches long an inch wide, petiole slender an inch long. Male panicle long and slender a foot or more, spikes slender one to three inches long, 3 or more in a whorl, rachis angled minutely pubescent. Flowers very small distant pubescent. Bracts shorter ovate. Sepals oblong ovate. Petals nearly as long oblong obtuse. Stamens 6, filaments fairly long.

Singapore on Bukit Timah (Ridley 4596.)

This wild yam is remarkable for its very thin leaves and slender stems. I have only met with a male plant but it seems very distinct from all other described species.

#### LILIACEÆ.

*Ophiopogon Malayanus*, n. sp. Stem erect stout 4 inches or more tall covered with the scarious sheaths of the fallen leaves and emitting long woody roots. Leaves linear acuminate.

Jour. Straits Branch

nate, falcate striate, glaucous beneath 9 inches long  $\frac{3}{4}$  inch wide. Scape 4 inches long base nude. Flowers small white. Bracts 2, bases broad ovate, tips linear, outer one  $\frac{1}{4}$  inch long. Pedicel longer. Sepals and petals ovate subacute  $\frac{1}{2}$  inch long. Stamens connivent filaments very short, anthers lanceolate. Style terete longer.

Perak at Padang Rengas (W. Fox) Lankawi on Gunong Raya (Curtis 2643). The *Ophiopogons* are abundant in the mountainous regions of North India and China, but get very scanty further south, in fact this is the only one from the Peninsula except the little known and doubtful *O. prolifera* from Penang. Though perhaps as near the common and variable *O. intermedius* of India, it is very distinct in its tall stout stem, broad leaves and ovate petals and sepals.

*Tupistra violacea*, n. sp. Terrestrial stem short and stout, leaves broadly oblong oblanceolate acute narrowed gradually to the base, 3 feet long, 3 inches wide, dark green stiff, spike very dense about 8 inches high. Flowers sessile  $\frac{1}{2}$  inch across, tube hemispheric violet, lobes oblong recurved darker. Anthers 6 sessile elliptic. Style cylindric white, spotted with violet. Stigma club-shaped rounded obscurely three-lobed.

Perak, Bujong Malacca; Penang, Highlands (Curtis).

This *Tupistra* differs from the other Peninsular species, *T. grandis*, Ridl., in its smaller lighter colored flowers and the rounded club-shaped stigma which is flat and circular in *T. grandis*. Though not perhaps as striking as that species it is a pretty plant.

*Dracaena conferta*, n. sp. Stem 4 to 10 feet tall, little or not branched. Leaves narrow linear acuminate 24 to 27 inches long  $\frac{3}{4}$  inch wide, base broadly sheathing, midrib very strong at the base, thinning upwards and disappearing before the tip. Raceme 2 feet long, erect, simple or occasionally branched, the base with numerous reduced leaves with broad bases, and acuminate upwards, peduncle rather stout with a few small sheathing leaves, inflorescence long dense. Flowers white, three or four in a tuft, with very short pedicels, and 2 or 3 ovate acuminate bracts  $\frac{1}{2}$

inch long. Perianth tube  $\frac{1}{4}$  inch long, narrow lobes linear  $\frac{1}{4}$  inch long. Fruit globose as large as a pea.

Perak, Selama at 300 to 500 feet elevation (King's collection 3149); Thaiping Hills near the Tea Gardens. There is also a specimen in Cantley's collections without locality, either from Malacca or Sungei Ujong. This *Dracaena* is allied to *D. Porteri*, Bak., but has longer and stiffer leaves with a very dense raceme of flowers. It is known to Malays as "Poko San Juan hutan jantan."

*Dr. penangensis*, n.sp. A shrub about 8 or 10 feet tall, with the stem an inch through, leaves when young oblong petioled, adults shorter. Upper leaves lanceolate acuminate 6 to 7 inches long, 1 to  $1\frac{1}{4}$  inch wide, petiole 1 inch long, sheathing at the base. Panicle erect 8 to 10 inches long with about eleven or twelve stiff branches four inches long or less with lanceolate acuminate bracts at the base  $\frac{1}{4}$  to  $\frac{1}{2}$  inch long. Flowers in twos and threes on slender pedicels  $\frac{1}{4}$  inch long. Perianth lobes narrow, white. Fruit as large as a cherry, 1 to 3 seeded, scarlet.

Penang at Batu Feringhi on the banks of the stream and at Highlands (Curtis 2369). Dried specimens of this often resemble those of *D. Mainyayi* our biggest tree *Dracaena*, but the living plant is much smaller and the leaves are very variable in form.

*D. Jackiana*, Wall. Cat. 5145. This plant obtained in Penang by Wallich was confused by Baker (Journ. Linn. Soc. XIV, 5-32) with *D. aurantiaca*, Wallich, under the name of *D. spicata* var. *aurantiaca*; and Hooker in the flora of British India puts it under imperfectly known species. I have not seen Wallich's specimens but Dr. Prain says that a plant collected on the Thaiping Hills by Kunstler (No. 2719) is exactly like the Wallichian plant. This plant is I think identical with a plant I collected in Pahang and described under the name of *D. longifolia* (Trans. Linn. Soc. III, 388). Hooker refers the Perak plant to Kurz's *D. pachyphylla*, an Andamanese plant, which however is probably distinct to judge from his description.

*D. pendula*, n.sp. A tall shrub with a moderately stout stem about 6 feet tall, leaves broadly oblong lanceolate acuminate.



inate narrowed to the base and subpetiolate. Petiole winged, 8 to 14 inches long  $1\frac{1}{2}$  to  $3\frac{1}{2}$  inches wide. Panicle graceful nodding 15 inches or more long with a few distant branches slender 7 or 8 inches long bracts small ovate lanceolate. Flowers in tufts of 2 or 3 distant white on pedicels  $\frac{1}{4}$  inch long, slender  $\frac{3}{4}$  to nearly an inch long split into segments for one-half their length; the lobes linear dilated upwards, stamens about as long, anthers oblong.

This fine plant grows in damp swampy spots in forests. I collected it in the Dindings on Gunong Tungal (No. 9448 of my collections) and have it also from Panchur in Malacca; and from Kwala Dipang and Gopeng from King's collections (Nos. 8279 and 4643). The Malacca specimens have broader and more distinctly petiolate leaves, and the panicle is stouter, but it appears to be specifically the same. It belongs to the *nutantes* section of *Dracaena*, in which the flowers are in long pendulous lax panicles.

*Dracaena elliptica*, Thunb. I found this common and variable plant growing abundantly in muddy swampy places along the Sungei Tebrau, this Easter. In this locality it had quite a different appearance from the common lowland dry forest form, being altogether a much larger and stouter plant, almost a small tree in fact.

We have now no less than sixteen species of *Dracaena* recorded from the Malay Peninsula, but there are doubtless more than this for I have seen several plants in our forests which appear quite distinct from any described, but of which I have not been able to obtain flowers.

#### COMMELINACEÆ.

*Forrestia gracilis*, n. sp. Stem creeping then ascending for about three feet a quarter of an inch through, twiggy dark green and glabrous, internodes 2 inches long terete. Leaves lanceolate acuminate narrowed into a winged petiole, dark green and glabrous above, velvety beneath, margined with red appressed hairs, 8 inches long 2 inches wide, petiole and mouths of sheaths hairy. Heads small

few-flowered; bracts ovate pubescent white with a green keel. Sepals oblong hooded ciliate white  $\frac{1}{8}$  inch long; petals longer lanceolate acute white; stamens 6; filaments contorted with a tuft of white hairs on the top. Anthers ovate deeply grooved white. Style filiform flexuous acuminate. Stigma minute, ovary small covered with white hair; capsule oblong pink.

*F. mollis*, Clarke, Monogr. Commel. p. 236 (in part, not of Hasskarl).

Very common in woods, Singapore. Tanglin, Pulau Ubin (Ridley 4130), (Walker 155); Johore: Tanjong Bunga (Ridley 6320); Pahang: Pulau Tijau (2381, 2382); Malacca: Sungei Buluh (10512); Selangor: Kuala Lumpur, Sungei Ujong: Gunong Berumbun, (Cantley's Coll.); Dindings: Telok Sera. Kedah Peak.

It is curious that this our commonest species has escaped being described till now, but it seems to have been mixed up with the rarer *F. mollis*. It is known as "Setawa betina" and the Sakais use a decoction of its roots for rheumatism.

*F. irritans*, n. sp. Stem tall and stout  $\frac{1}{2}$  inch through, leaves elliptic lanceolate with a broad flat petiole, acuminate 9 inches long, 3 inches across, hairy on both sides, sheaths  $1\frac{1}{2}$  inches long, glabrous except for the hispid edges. Heads large and dense over an inch through, densely covered with red spiny hairs, sepals lanceolate covered with similar hairs. Stamens 6, anthers oblong, style filiform.

Selangor, on the Tras route at the 15th mile; Perak Bujong Malacca (Ridley 9784), Sungei Ujong, Bukit Kupayiang (Cantley's collection.) Rather rare in woods at an altitude of about 2,000 feet. A very distinct species, from its large globose heads of flowers covered densely with sharp pungent red spines. I have never seen fruit of it.

#### PALMÆ.

*Pinanga Singaporensis*, n. sp. Stems tufted dark green 6 to 15 feet tall  $\frac{3}{4}$  inch through; internodes 8 inches long.

Johr. Straits Branch

Leaves concolorous or a little paler along the back about 3 feet long, sheath tubular, petiole 8 inches long, leaflets sigmoid acuminate about 13, sub-opposite, terminal one forked; 12 inches long by 3 inches wide or less, nerves 5 to 7. Spike branched erect 4 inches long, with 5 branches. Spathe thin boat-shaped papery with a short point brownish pink  $1\frac{1}{2}$  inch long. Rachis cream color scurfy, flowers distichous. Males  $\frac{1}{4}$  inch long, ivory white. Sepals very small acute. Petals oblique ovate acuminate much larger. Stamens 14 white, filaments very short, anthers oblong, no pistillode. Female petals and sepals broadly oblong pink, edges ciliate, stigma small, sessile papillose circular white.

This pretty palm occurs in Singapore in forests at Bukit Timah, Mandai, Stagmount and Selitar, and is No. 11267 of my collections. I have not seen it outside Singapore, and it is rather scarce there.

*Ptychoraphis longiflora*, n. sp. Habit and stems of *Pt. Singaporensis*.

Leaf three feet or more long, rachis brown scurfy, leaflets linear acuminate 5-ribbed, alternate about 40 pairs 18 inches long,  $\frac{1}{2}$  inch wide. Peduncle stout 1 inch long ribbed and scurfy, branches of inflorescence slender 4 or 5 scurfy, 6 or 7 inches long. Flowers in distinct pairs, one male and one female together about 20 pairs on a branch; males  $\frac{3}{8}$  inch long, with two orbicular gibbous bracts at the base. Calyx lobes orbicular fringed. Petals lanceolate obtuse much longer. Stamens 6 shorter than the petals, filaments with a broad base narrowed upwards, anthers oblong dorsifixed. Pistillode conic stout as long as the stamens. Female flowers. Bract single small. Sepals orbicular ciliate. Petals ovate orbicular imbricate, pistil ovoid.

Johore on the top of Gunong Banang at Batu Pahat, (Ridley 1121). Besides this there are two other species of this genus described. *Pt. Singaporensis*, Becc., the Korintin palm, abundant in the South of the Peninsula, and *Pt. angusta* of the Nicobars, a plant of totally different habit. This new species closely resembles *Pt. Singaporensis*, but the petals of the male are very much longer.



*Iguanura spectabilis*, n. sp. Stem from 3 to 8 feet tall, tufted or solitary very stout over  $1\frac{1}{2}$  inch through. Leaves entire and cuneate bilobed at the apex, margins crenulate toothed strongly ribbed, three or more feet long one foot across, petiole. Inflorescence from among the leaves, peduncle over a foot long with two sheaths 6 inches long, and a bract-like ovate acute leaf  $\frac{1}{4}$  inch long above, branches stout ascending, six inches long, deep green. Flowers spirally arranged sunk in the rachis. Male, sepals ovate. Petals oblong acute twice as long  $\frac{1}{8}$  inch long. Stamens 6, longer than the petals, filaments thick connate at the base with the long cylindric obtuse pistillode. Females, sepals rounded, petals ovate rounded. Fruit elliptic tip curved, black when ripe,  $\frac{3}{4}$  inch long, on the thickened rachis.

This superb palm known to Malays as the "Teruno" occurs on the Hermitage Hill in Perak, and at Bruas in the Dindings where it was first obtained by Mr. Curtis and later by myself, (No. 8403) of my collections. Young living plants sent home by Mr. Curtis were described by Dr. Masters in the Gardener's Chronicle (1898, XXIII, 258) under the name of *Geonoma Pynaertiana*. In the Dindings it forms a bush with numerous short stems, on the Hermitage Hill I found it with a stout tall stem. It is a most beautiful palm when well grown, the broad leaves deep green when adult, and pink when first put out, making it most attractive.

*I. ferruginea*, n. sp. Stem slender, leaves 18 inches long, petiole four inches long, rachis red-scurfy, leaflets 5 or 6 pairs alternate trapezoid, base narrowed, apex acuminate, margins more or less toothed, 6 inches long 3 inches through. Spathes 6 inches long acuminate. Inflorescence on a peduncle 6 to 8 inches long, red tomentose, branches thick 9 to 11 some-times branched again, 6 inches long covered with red wool. Flowers scattered distichous in foveolæ, covered with red wool. Male flowers glabrous. Sepals ovate rounded, petals lanceolate ribbed,  $\frac{1}{3}$  longer. Female flowers sepals similar, petals shorter and

blunter. Drupe cylindric curved, slightly narrowed towards the tip, black, half inch long. Seed cylindric acute curved rugose  $\frac{3}{8}$  inch long; albumen equable.

On the Thaiping Hills common from about 2000 feet elevation upwards (10684; 11405 of my collection) and Bujong Malacca (Curtis 3164). This palm has the habit of *I. polymorpha*, Becc., but the stouter inflorescence on a longer peduncle and covered with red wool, the larger flowers, and curved cylindric fruit, make it very distinct.

The Iguanuras are not an easy group of palms to separate, as they are apt to be very variable especially in the form of the inflorescence. There appear to be seven kinds at present known from the Peninsula. The commonest species is *I. geonomæformis*, Mart. A lowland species with usually one simple spike. Of this there are the following forms: (a) *typica*, with a simple stout tomentose spike, (b) *ramosa*, a similar spike but branched, with 2 or 3 branches, (c) *malaccensis*, with a slender glabrous simple spike (occasionally branched). The leaves of all these forms may be quite entire. *I. Wallichiana*, Hook fil., with a compound inflorescence of several slender branches, is allied to this latter form. It also varies in the form of the leaves. Two plants described by Hooker and Beccari from Scortechini's collection, viz., *I. diffusa* and *I. parvula*, appear to me to be forms of these species. The first is a large plant with an inflorescence more compound, the second a dwarf abnormal form. I have only however seen sketches of them and very little of the plants appears to have been collected. *I. corniculata* has also only once been collected. The fruit is curiously curved, and is possibly abnormal. *I. polymorpha*, Becc., including *I. brevipes*, Hook. fil., is a distinct plant, not rare on the Thaiping Hills. *I. ferruginea* is allied to it but seems to me to be distinct. *I. bicornis* has curiously bilobed fruit, and occurs also on the Thaiping Hills, but seems to be rare.

*Livistona rupicola*, n. sp. Stem short and thick about 3 feet long and 6 inches through, densely covered with brown fibers from the leaf sheaths. Leaves orbicular about 2 feet across with about 12 leaflets rather narrow very acumi-

nate tips setaceous; petiole slender 20 inches long nearly thornless. Spadix slender, much branched. Spathes split striate cuspidate dark brown, the two basal ones three inches long, upper ones slightly dilate cuspidate. Branches many long slender with many very long and slender sinuous spikes 1 to 2 inches long, lengthening to 4 inches in fruit. Bracts at the base of the branches linear narrow caducous. Flowers very small globose; sepals rounded gibbous. Petals similar. Staminal crown with 6 subulate teeth.

Selangor limestone rocks at the Kuala Lumpor caves (8285 of my collection), Lankawi Islands, (Curtis).

This charming palm, the Serdang Batu of the Lankawi Islands, is probably the smallest species of the genus. It is remarkable for its very short thick stem densely covered with brown fibres, the remains of old leaf sheaths. The inflorescence is also very small for the genus. It grows on the limestone rocks of the Selangor caves from the lower part to the top of the cliffs.

*Licuala longepedunculata*, n. sp. Stem 2 to 8 feet tall, 1 to  $1\frac{1}{2}$  inch through, leaves 4 to 5 feet long orbicular, lobes 6 or more narrowed at the base and broadened upwards with broad blunt teeth 15 to 16 inches long and 3 inches across at the tip, deep green, petiole slender not thorny. Inflorescence slender 3 feet or more long, peduncle broad flattened at the base, quite glabrous over 2 feet long, sheaths distant long, spikes 3 inches long, in fours. Flowers scattered over  $\frac{1}{4}$  inch long, rachis moderately stout, calyx urn-shaped sessile not lobed. Petals lanceolate obtuse.

Perak, Gunong Batu Putih (Wray 254) (King's Coll. 8148). I only know this species from the above mentioned collections. Beccari labels it "*L. glabra*, foliis latioribus," but it is certainly very distinct from *L. glabra*.

*L. (Pericycla) paniculata*, n. sp. Leaf large, lobes  $2\frac{1}{2}$  feet long, rather narrow, with short teeth  $\frac{1}{4}$  inch broad. Inflorescence paniced wide-spreading 3 feet long, spathes 6 not split, the lowest 8 inches long  $\frac{1}{2}$  inch wide, upper part reddish, upper spathes shorter dilated upwards. Branches



panicled, spikes long and slender 5 to 7, glabrous or scurfy, 7 inches long or less. Flowers scattered sessile  $\frac{1}{8}$  inch long. Calyx cylindric with a broad base, truncate obscurely and irregularly lobed glabrous. Petals short and broad ovate acute. Staminal ring with 6 teeth. Pistil sometimes bilobed, stigma lateral.

I collected a specimen of this on the Hermitage Hill in Perak, and have never seen it since, and unfortunately omitted to record the size of the plant. It is evidently allied to *L. pericycla*, Zipp., *Pericycla penduliflora* of Blume, a native of New Guinea, from which it differs in having solitary sessile, not stalked flowers in pairs. The panicled inflorescence is unique in Malay Licualas as at present known, and is the characteristic of the section *Pericyclus* of which the New Guinea species is the only other one known. It is to be hoped that any one visiting the Hermitage Hill will recollect this curious plant and bring full details of its habit and size.

*Calamus aquatilis*, n. sp. Stem fairly stout about 30 feet long covered with dense bristle-like black spines. Leaf large flagelliferous armed with copious black spines in short rows, petiole a foot long stout black rounded, edge armed with spines of different sizes flattened 2 inches long or less, those at the base black long or slender, leaflets numerous equidistant linear acuminate 1 foot long an inch broad, bristles short scattered along the edges and back, flagellum 4 feet long, hooks in half whorls numerous. Panicle very large and stout, lower sheaths tubular an inch through with numerous decurved spines single or in twos and threes black tipped, chiefly on the back and sides on the upper part. Branches about 2 feet long, secondary spathes tubular unarmed an inch long, male branchlets numerous 6 inches long, spathels tubular about  $\frac{1}{2}$  inch long, spikes subdistichous recurved  $\frac{3}{8}$  inch long green. Spathellules saucer-shaped with a small ovate limb, sepals oblong lanceolate. Petals as long oblong lanceolate thicker. Stamens, filaments rather long slender, anthers long narrow. Female branchlets 2 to 3 inches long, spathels funnel-shaped. Spathellules

small saucer-shaped. Flowers solitary, sepals ovate as long as the petals and quite similar. Fruit small globular beaked  $\frac{1}{2}$  inch long scales in 6 whorls, rhomboidal as broad as long, yellow or brownish yellow, tips darker, grooved.

This rattan is common in tidal river swamps, and is known as "Rotan Bakau." There is a trade rattan of the same name but I am not yet certain that it belongs strictly to this plant. The general appearance of this species when out of flower is that of *Daemonorops angustifolius*. It does not seem to flower very frequently, as I have seldom found it in flower or fruit, abundant as it is, which is probably why it has never hitherto been described. Professor Beccari who is engaged on a work on our rattans to be published in the Annals of the Botanic Gardens of Calcutta, considers it to be quite a distinct plant from any described.

I have met with it in Singapore, on the Serangoon Road, and at Changi (6275 of my collections), Johore, Panchur on the Johore river; on the Sungei Tebrau river, and at Batu Pahat (11216), and in Pahang at Kuala Pahang.

#### AROIDEÆ.

*Cryptocoryne purpurea*, n. sp. Aquatic, stem creeping stoloniferous.

Leaves ovate or elliptic ovate, base broad 3 inches long  $1\frac{3}{4}$  inch wide, petioles 4 inches long, peduncle short about half an inch long, spathe 3 or more inches long, tube twisted  $\frac{1}{4}$  inch through, white, limb ovate caudate  $\frac{1}{2}$  to 1 inch long, half an inch wide deep purple brown with a yellow mouth, pustular, tail  $\frac{1}{2}$  inch long, valve over the inflorescence oblong truncate pale yellow. Female flowers 6 to 8 connate in a circle, styles very short, stigmas discoid, neuters above the pistils 5 or more, nude portion of spadix slender. Male flowers few yellow obcuneate, appendix small elliptic clubbed.

Johore, Kota Tinggi (4214 of my collections). This plant cultivated in the Singapore Botanic Gardens for several years, was sent to Kew Gardens where it flowered

and was figured in the Botanical Magazine plate 7719 under the name of *C. Griffithii*. It is however quite distinct from that in its more numerous, female flowers, smaller appendix to the spadix, and much longer tube. It grows very readily in water, and is perhaps the easiest to cultivate. I have only seen it in the one locality mentioned, our commonest species being the short tubed *C. Griffithii*.

*Typhonium fultum*, n. sp. A small herb with a short stem above the ground propped on strong roots. Leaves 5 or 6 ovate or deltoid hastate 2 to 3 inches long 2 inches wide deeply cordate, apex acute, lobes rounded, petiole  $2\frac{1}{2}$  inches long. Peduncle  $\frac{1}{4}$  inch long, spathe with a green base dilated  $\frac{1}{4}$  to  $\frac{1}{2}$  inch long, limb 2 inches long  $\frac{1}{2}$  inch broad; subacute or acuminate purple-veined, tip green. Spadix about as long. Appendage orange brown cylindric from a thick base tapering upwards, below a short nude portion. Male portion of about 6 rows of oblong yellowish anthers, then a longer nude portion. Neuters numerous horn-shaped white up-curved. Females in one whorl oblong, one ovuled.

Selangor at the limestone caves near Kuala Lumpor (8165) also found in Lankawi Islands by Curtis.

This odd little aroid is remarkable for the stem standing up supported on its roots above ground after the manner of a Pandanus, a habit not observed in any other species, the rest of them having small underground tubers.

*T. filiforme*, n. sp. A very small herb, tuber globose  $\frac{1}{2}$  inch through. Leaves 2 or 3 deltoid hastate acuminate, lobes divaricate acute 2 or 3 inches long,  $\frac{3}{4}$  to  $1\frac{1}{2}$  inch wide, petiole slender 2 to 5 inches long. Peduncle  $\frac{1}{4}$  inch long. Spathe 3 inches, base elliptic  $\frac{1}{4}$  inch long olive green, limb linear very narrow, apex eventually coiled up yellowish spotted with red. Spadix very slender  $3\frac{1}{2}$  inches long, appendix ochre yellow filiform nearly 3 inches long, male portion short below them a long slender nude portion, neuters filiform contorted long, female flowers few oblong.



I found this at the foot of the limestone cliffs at Kuala Dipang in Perak. It is evidently allied to *T. bulbiferum*, Dalz., of the Concan, differing in the absence of any trace of bulbils, the longer appendage and neuters, and longer bare space below the male flowers.

*Amorphophallus Malaccensis*, n. sp. Tuber 4 inches through hemispheric with a depressed top. Leaves not certainly known. Peduncle 7 inches tall rough. Spathe tube wide trumpet-shaped 3 inches long, limb broad rounded six inches long and as wide. Spadix longer, appendage fusiform acuminate 6 inches long rugose hollow, loose textured and fibrous within. Male portion one inch long anthers crowded oblong. Female portion  $2\frac{1}{2}$  inches long. Flowers numerous, ovaries ovoid, style  $\frac{1}{4}$  inch long.

I only know this from some dried specimens which were collected by a native collector on Bukit Panchor in Malacca. It belongs to the section including *Amorphophallus*, *Rex*, and *campanulatus*, but is distinct in its longer peduncles, and longer narrower appendage. I have several times met with foliage of a plant very much resembling that of a *Rex* with a tall stout dark green scabrid petiole which may perhaps belong to this species. A very large spike of fruit over a foot longer with berries half an inch long which was obtained by Mr. Hervey on Bukit Payong in Malacca is very probably the fruit of this plant. It should be looked for again.

*A. minor*, n. sp. Tuber an inch through. Leaf petiole slender 9 inches long, blade three branched each branched again, leaflets numerous thin green (red when young) lanceolate acuminate inaequilateral 3 inches long by one inch wide, nerves numerous close joining an inner intramarginal one. Peduncle 8 to 12 inches long with loose brown sheaths at the base, the largest truncate 4 inches long. Spathe 3 to 4 inches long convolute at the base lanceolate acuminate 1 to  $1\frac{1}{2}$  inch across. Spadix 3 inches long. Appendage cylindric-conic an inch long. Male portion as long as the appendage. Flowers very numerous crowded. Females few. Style very short.

Perak on the Thaiping Hills; and in Province Wellesley at Tasek Gelugor. This is allied to *A. sparsiflorus*, Hook. fil., differing in the shorter appendage and longer male portion of the inflorescence.

*A. carnea* n. sp. Tuber hemispheric 3 inches across. Leaf unknown. Peduncle 3 inches and a half long, smooth gray green with brown blotches, sheath at base 2 inches long. Spathe 5 inches long, loosely convolute at the base with a broad limb pinkish with brown blotches, paler within. Spadix rather longer. Appendix conic obtuse 3 inches long 1 inch through at the base fleshed low obscurely cancellate. Male portion 2 inches long stout greenish. Stamens densely crowded. Female portion ovaries shortly pedicellate, broad flask shaped, style short and thick. Stigma lobed.

Lankawi Islands (Curtis). This plant was flowered from a tuber brought in with tubers of *Arisaema* from Lankawi Islands, in the Penang Gardens. I have only seen a drawing of it but I know nothing quite like it.

*Alocasia ovalifolia*, n. sp. Stem about a foot tall, leaves ovate cordate acute dark green one to 2 feet long 5 to 12 inches wide with 20 pairs of nerves, petiole stout over a foot long. Peduncle six inches long or less. Spathe 3 to 6 inches long, tubular portion  $1\frac{1}{2}$  inch long, limb 3 inches long  $\frac{1}{2}$  inch wide oblong obtuse. Spadix slender, appendage cylindric  $\frac{3}{4}$  to 2 inches long. Male portion 1 inch long. Flowers oblong crowded, below them a nude portion. Female portion 1 inch long.

Johore, base of Gunong Pantu; Selangor, Tras Route at the 15th mile (No. 8487), Ginting Peras, Bukit Kuta; Perak, Sungei Larut (Wray 2457), Larut Hills; Penang, Moniots Road (Curtis). Rather a dull looking *Alocasia* not uncommon in the hill woods.

*Pothos inaequalis*, n. sp. Stem rather slender much branched. Leaves elliptic oblanceolate with a long point and acuminate base intramarginal vein often far from the edge undulate with another very inconspicuous one close to the edge, main nerves often large and remote 4 to 7

inches long 1 to 2 inches wide. Petiole  $\frac{1}{2}$  to 1 inch long sheathing to the thick knee which is half an inch long. Bracts several, upper one sheathing  $\frac{1}{2}$  inch long, lower ones small. Peduncle  $1\frac{1}{2}$  to 2 inches long fairly stout curved. Spadix an inch long cylindric dense flowered. Sepals cuneate with a broad flat triangular top. Pistil top broad rounded. Pahang, Pulau Tawar (2391).

*P. ellipticus*, n. sp. Stem fairly stout leaves elliptic cuspidate 5 inches long  $2\frac{1}{2}$  inch wide, pale green when dry, keel stout, base rounded, inner intermarginal vein some way from the edge, outer one close to the edge, parallel nerves fine and close. Petiole 3 inches long sheathing rather broadly to the knee  $\frac{1}{4}$  inch long. Peduncle stout 2 inches or less deflexed. Spadix nearly sessile  $\frac{3}{4}$  to 1 inch long thick cylindric dense flowered. Flowers fairly large. Sepals oblong with a broad inflexed top. Stigma discoid shortly elevated. Fruit oblong  $\frac{1}{2}$  inch long.

Pahang on the Pahang river, at Kuala Tembling and Pulau Tawar.

*P. grandispatha*, n. sp. Stem slender less than  $\frac{1}{2}$  inch thick. Leaves lanceolate cuspidate 6 to 8 inches long, 2 to  $2\frac{1}{2}$  inches wide, cusp  $\frac{1}{2}$  inch long base narrowed blunt. Petiole 2 inches long sheathing rather broadly to the very short  $\frac{1}{2}$  inch knee, sheaths eventually breaking up into fibres. Peduncle slender 2 inches long. Spathe broadly ovate acuminate, apparently purple in life, narrowed at the base  $2\frac{1}{2}$  inches long 1 inch wide. Spadix very slender 1 inch long sessile. Flowers very small, sepals oblong truncate, style distinct with a small discoid stigma.

Penang, West Hill (Curtis). Allied to *P. brevistylus*, Engler. This plant is remarkable for its large spathe.

#### PANDANACEÆ.

*Pandanus immersus*, n. sp. A stout pandan growing immersed in water, the stem rising but little above, leaves many feet long broadly linear 4 inches across, glaucous beneath, acuminate with a long point, thorns numerous brown. Female spike solitary on a stout flexuous peduncle 2 inches long  $\frac{1}{2}$  inch through, oblong thick green 4 inches long  $1\frac{1}{2}$



inches through, carpels small with a very stout simple curved style  $\frac{1}{8}$  inch long.

Selangor in the Labu River, forming dense thickets.

Native name "Mengkuang Ayer."

- P. bicornis*, n. sp. Stems short, only one or two feet tall. Leaves dark green glaucescent at the top of the stem 2 feet or more long 2 inches wide broadly linear and tapering quickly into a very thorny tail 3 inches long, edge and keel armed especially at the base with numerous recurved thorns. Female bracts broad ovate, outer ones rather abruptly acuminate 6 inches long with a slender point 2 inches long thorny, inner bracts obtuse, head solitary oblong rounded at the top 4 inches long and nearly as broad green, carpels nearly  $\frac{1}{2}$  inch through  $\frac{3}{4}$  inch long top broad free obscurely angled. Style broad short with two spreading short sharp horns with broad bases about  $\frac{1}{8}$  inch long.

Perak, Taiping Hills in dense forest at 2500 to 3000 feet altitude.

- P. aurantiacus*, n. sp. A large branching shrub with stems 2 or 3 inches through and about 12 feet tall. Leaves over 3 feet long,  $1\frac{1}{2}$  inch wide, glaucous green, linear long acuminate, tips deflexed, with numerous close-set short thorns on the edge and keel. Female inflorescence with a stout rachis about a foot long and five globose oblong heads  $2\frac{1}{2}$  inches long 1 inch through, orange colored. Carpels large  $\frac{3}{4}$  inch long, apex conic ending in a long stout but not very hard beak.

Singapore, Bukit Mandai Road; Johore, Tanjong Bunga (6288 of my collection); Perak, Larut Hills; also in Sarawak.

This grows on swamps on river banks usually near the sea, and is known in Johore as "Pandan Akar". This may possibly be the *P. Yvoni* of Solms-Laubach (*Linnaea* XLII, 20) based on a plant collected by Yvan in Malacca, but the description is too poor to identify any pandan.

- P. stelliger*, n. sp. A slender stemmed pandan 4 to 10 feet tall, branched. Leaves linear acuminate shining grey green

over a foot long and an inch wide, the point narrow 3 inches long, thorns distant except at the point which is very thorny. Heads oblong ovoid 2 inches long and over an inch through in a cluster of four nearly sessile. Carpels  $\frac{1}{2}$  inch through the style oblong, top square with from 3 to 6 horizontal points sharp and rough at the tip.

Selangor, on the Tras Route at the 20th mile (8775;) Perak, Thaiping Hills at 2500 feet altitude in forests.

This was referred to *P. minor*, Ham., by the Kew authorities but differs in the clustered heads of fruits and the curious style which is quite flat and square at the top with from 3 to 6 horizontally spreading points.

*P. glaucophyllus*, n. sp. Stems 10 to 12 feet tall  $\frac{1}{2}$  to 1 inch through. Leaves crowded at the top 1 to 2 feet or more long 2 inches wide glaucous broadly linear oblong, tip abruptly acuminate, edge minutely denticulate rather flaccid, midrib prominent with very small thorns, capitulum globose 2 inches long on a stout peduncle about 3 inches long. Fruit rounded tapering into a stout upcurved simple spine  $\frac{1}{6}$  inch long grooved above the tip acute.

Perak, Thaiping Hills in dense forests, about 2500 feet altitude. Rather local, growing in small patches. Allied to *P. parvus*, Ridl, but very much larger in all its parts.

*P. penangensis*, n. sp. A tall stout tree about 20 feet tall and four inches through the stem. Leaves very long linear acuminate 6 to 12 or 14 feet long 4 inches across often especially when young marbled light and dark green, edges and keel armed with stout thorns crowded towards the tip of the leaves. Heads 4 or 5 together on a short peduncle, green oblong 6 inches long, 3 inches through. Carpels an inch long, free portion conic angled  $\frac{1}{3}$  inch long. Styles strong simple hard dark brown polished curved forward  $\frac{1}{2}$  an inch long.

Penang Hill from the base to the top, in woods. This resembles *P. furcatus*, Roxb., in habit but has a compound inflorescence.

## CYPERACEÆ.

*Mapania triquetra*, n. sp. Stem ascending or erect 6 inches tall, trigonous broad from the very broad leaf. Sheaths 12 inch long and one inch wide green edged with pink. Petiole green 2 to 3 inches long, blade oblong dark green above, paler beneath, base broad apex rounded ending abruptly in a tail  $1\frac{1}{2}$  inch long, edges aculeate, whole blade 6 to 8 inches long, 2 inches wide. Scapes slender 3 inches long red. Head of flowers very small  $\frac{1}{2}$  inch long narrow. Bracts shorter lanceolate red. Squamellæ 6, two outer ones with ciliate keels. Stamens three white anthers twisted linear oblong. Style long. Stigmas 3 short.

Woods, Dindings, Lumut (Ridley), Pangkor (Scolotechnii); Negri Sembilan, Bukit Tumiang (Cantley's collection); Penang Hill (Curtis). Also in Sarawak.

This pretty and curious plant has been confused with the common *M. humilis*, Naves and Villar, but is really totally distinct in the broad triangular stem shorter and broader leaf blade and the very small head of flowers.





## Notes on a Cruise in the Southern China Sea.

C. BODEN KLOSS.

In 1900 I spent about eleven weeks, including the months of August and September, cruising about with Dr. W. L. Abbott in his Schooner "Terrapin" which had just been launched. Our purpose was to make collections of mammals and birds and of any other objects zoological that might fall in our way. As the islands of the Southern China Sea amongst which the time was spent, have received either but scanty notice or none at all, the following pages may have some interest. Unfortunately for the present purpose I kept only the very baldest log of our voyage so that the account of our experiences, drawn up after an interval of three years, is far from being as satisfactory as I could wish.

I was unable to accompany the schooner when she sailed at the beginning of July so arranged to join her at Linga, and in the middle of the month therefore left Singapore in the *S. S. Malacca*.

We stopped a night at Rhio *en route* and I was thus enabled to get from the Resident a permit to travel in the Rhio-Linga Dependency in which the whole of our cruise was to take place. The town of Rhio is prettily situated and laid out, but very small and quiet; it possesses an hotel and a good fruit-market; every other one of its shops appears to be run as a pawn-broking establishment. A long walk in the surrounding country showed me nothing more inviting than clayey hills covered with bracken and the *S'ndudu* tree (*Melastoma polyanthum*), and everywhere dotted with Chinese tombs.

From Rhio to Linga—we called at Sinkep Island on the way with provisions for the tin mines—was, I think, four days steaming through smooth seas and green islands. The *Macassar* is an old wooden tub capable of doing about 7 knots in calm water with the wind astern. The only accommodation

she could offer was the open bridge crowded with natives, where I found room for a deck chair. Had this been all there would have been nothing to kick at, but unfortunately the vessel swarmed with a certain highly objectionable and active *Rhynchota* that spoilt all pleasure, and when I hailed with delight the appearance of the "Terrapin" lying off Tanjong Buton on the south side of Linga, I was, after four days of an inferno, in a state that I had never been in before and fervently pray never to attain again.

#### Linga.

Lying about mid-way between Singapore and Banka, Linga is an island of irregular shape about 33 miles in length in a north-west and south-east direction, and is surrounded by smaller islands of various sizes. Unlike its near neighbour Sinkep, it is not worked for tin, and is best known as being at one time the head quarters of the numerous pirates who used to ravage the western seas of the Malay Archipelago in the early part of last century.

The schooner had to anchor a mile or so from the land since off-shore for some distance extended banks of soft black mud through which we were compelled to wade when the tide was low while at other times the sea broke on them with some force for the roadstead is exposed to south-easterly winds. At Tanjong Buton were a few Chinese *kedais* and the house of a Dutch Assistant-Resident (now withdrawn) and from here a road had been made to the town of Linga.

The best collecting ground was on the outskirts of the village of Maruang, lying two or three miles away between the road and the sea. The surrounding country was, for the most part, a sago swamp, but in the fruit plantations of the Kampong certain birds and small mammals were numerous. The village itself consisted of a settlement of Sumatran Malays, the houses—about forty in number—built in two orderly rows with the mosque in the centre. The thousands of huge durian trees that surround it, were just then fruiting and at the little watch-houses in the plantations freshly fallen durians could be had in piles at a cent or two apiece while the few small steamers that call were constantly taking cargoes up to Singapore.



Learning from the people of the village that flocks of fruit-bats visited their fruit-trees nightly, we, one evening when there was a little moonlight, went up the plantation for the purpose of obtaining specimens.

Accompanied by half the boys of the Kampong, we took up places beneath a huge rambutan tree—the gathering point of great numbers of bats—but for a time made very poor practice at the dark bodies flitting between the branches in the dim light. Swarms of mosquitoes did not help to improve matters. At length however I hit on a plan that gave better results; choosing a large bunch of fruit that was frequently visited, I rested my gun against a convenient tree-trunk and took careful aim at it. Holding the barrels steadily in position all I had now to do was to stand up and watch the target: whenever a bat settled on that particular bunch I pressed the trigger and the shot was invariably followed by a thud on the ground or crashing and squawking among the branches as a wounded animal dropped slowly downwards. In a couple of hours the boys had picked up a dozen specimens of *Pteropus vampyrus* with which we returned to the boat, leaving sundry others to be recovered on the morrow.

On the way back a bullet was kept in readiness for wild-pig. None were seen, however, though nightly rootings along the road showed their presence in the neighbourhood.

The following day was given over to the preparation of skins and skeletons from the specimens obtained.

The 24th of July, our last day in the island, was spent in visiting Linga Kampong to bid farewell to the Sultan and to buy supplies for the voyage ahead.

The town is distant about an hour and a half from the sea and is reached by a road constructed by the Dutch Assistant-Resident. Although roughly made it is passable for the Sultan's gharry and for the few local rickshaws which, old and dilapidated, generally traverse it at a walking pace.

For the first two miles it runs through a swamp planted with sago palms, then coming out on hard clay ground is bordered with scrub and lallang until near the town it passes through plantations of various fruit-trees, sugar-cane and bananas.

Although the actual control is in the hands of the Dutch from whom he receives an income of some \$80,000 a year, the Sultan of Linga is nominally ruler over neighbouring parts of Sumatra, all the islands between Sinkep and Singapore and all the various small groups in the Southern China Sea.

The town of Linga—his capital—is situated on the banks of a stream navigable by small praus at high water, about two miles from its mouth. To the north a fantastically split peak, the highest summit of the Linga group, rises to a height of 3,921 feet, densely covered with jungle and scrub and scarcely ever free from clouds.

The population of the town, Malays and Chinese, number about 6,000. All the houses are built on piles, those of the Chinese on a muddy expanse by the banks of the river which not infrequently overflows. A number of substantial brick buildings have at one time also been erected by them but are now in a ruinous condition.

The houses of the Malays, shaded by fruit trees, arecas and coco-palms were scattered about without regard to orderly arrangement. Amongst them stood the school, which was apparently well patronised, and the Sultan's palace, a large and ugly barn-like structure of wood.

The Sultan was absent and we proceeded at once to the Chinese portion of the town to get through with our marketing.

The Chinese community is under a "Captain China" who in the Linga Sultanate is responsible to the Dutch only for the good behaviour of his charge and who collects for them the poll tax of \$3 a head every year.

The bazaar was of fair size and sold the usual merchandise found in such places that includes various articles ranging from a bottle of scent to an onion. Our requirements of rice, vegetables, curry stuffs etc., were soon satisfied and from the sarong shops kept by Klings we got a few European cotton sarongs for use on board. The Malays here were apparently like the lillies of the field. "They toil not neither do they spin," and it was with difficulty that we got from them even a few eggs and chickens.

We returned to the "Terrapin" by way of the river in a canoe with a small roof of kajangs. The water was very low

and paddling between muddy banks we were unable to see anything of the country on either side. Crocodiles are said to be numerous but none were met and after crossing the bar at the river mouth, we got up sail and in short time reached the schooner.

Previous to my arrival Dr. Abbott had made an expedition to the peak. He spent eight days in a hut built at an altitude of about 1000 feet and ascended on three occasions to 3000 feet, but each time the mist was so dense that he could not proceed higher. Animal life was remarkably scarce in the mountain forest; the lotong, however, was seen occasionally and at the highest elevation reached the "house" of a pig was found. Birds were very few in number. The mountain seemed to offer most attraction to a botanist: orchids occurred in great abundance.

In order to give a complete list of the mammals known from Linga I have included here the further material obtained by Dr. Abbott during a second visit to the island in 1901. All species that were described for the first time from both collections are distinguished by the addition of *sp. nov.*

1. *Semnopithecus maurus.*
2. *Macacus cynomolgus.*
3. *Pteropus vampyrus.*
4. *Tupaia tana.*
5. *T. malaccana.*
6. *Viverra tangalunga.*
7. *Arctogalidia simplex, sp. nov.*
8. *Tragulus javanicus.*
9. *T. pretiosus, sp. nov.*
10. *Sus vittatus.*
11. *Ratufa notabilis, sp. nov.*
12. *Sciurus vittatus.*
13. *Sciurus tenuis.*
14. *S. notatus.*
15. *Rhinosciurus laticaudatus*
16. *Mus lingensis, sp. nov.*
17. *M. fremens, sp. nov.*
18. *M. firmus, sp. nov.*



The birds of Linga show no peculiarities and do not differ from those occurring in the adjacent mainlands. The most complete collection recorded is that made by the native hunters of the late Mr. A. H. Everett. The list of species is given in "Novitates Zoologicae."

#### Pulo Taya and the Nyamok Islets.

We left Linga at 2 a.m. on the morning of July 25th bound for a group of three small islands forty miles to the south-eastward. With the wind ahead all the way it was not until mid-day on the 26th that we anchored off Taya the largest of the three.

The island is fairly steep-to, of granite formation rising to a double peak about 600 feet high. It is about a mile and a half long north and south, oval in shape and covered with forest.

Landing on the east side on a sandy bay in a bight between the hills we found at the south end of the beach just within the jungle a spring of good water. Near the shore the bay was blocked up with coral over which at low tide we had to scramble. A little party of Orang Laut in their crazy praus, visiting the island for "ikan merah" for which it is well known, told us that rats, squirrels, and a "biawak" were to be got; but during the three occasions we were ashore we saw neither. The only birds obtained were the Nicobar (*Calonotus nicobarica*) and Nutmeg pigeons (*Myristicivora bicolor*), which last occurs on nearly all islands in this region: the glossy starling (*Calonotus chalybeus*) with dark metallic green plumage and red irrides; a gaudy little sunbird (*Anthothreptes rhodolaema*), the Eastern reef heron (*Leptorodius sacer*), and *Halcyon chloris*, the blue-and-white kingfisher. Last and best of all was one specimen of *Columba grisea*, a bird of extreme rarity in collections.\* In general appearance it is somewhat similar to the cream-and-black Nutmeg pigeon if the former colour were replaced by a pale grey.

On the Nyamok Islands about a mile north of Taya, two islets, the larger no more than two or three hundred yards in

\* The British Museum Catalogue and Pigeons records a single specimen only.

diameter, we shot a blue-and-white king fisher and the reef heron.

These islets, neither of which is more than 150 feet high, are covered with thin jungle in which we saw several specimens of the Nicobar pigeon—the “burong mis” or golden bird of the Malays.

Taya was left at midday on the 28th and soon after making sail a squall from the south-west struck us. We ran before it, goosewinged with scandalised sails, the seas racing up behind and breaking in showers of spray under the counter. In the couple of hours it lasted we had made nearly twenty miles of our way to Pulo Pengiki Besar and afterwards sailing with a wind that allowed an easy course to be laid, anchor was dropped in a bay on the north side of the island at six o'clock on the evening of July 31st.

#### Pulo Pengiki Besar or St. Barbe Island.

Seen from a distance Pengiki appears like two or three separate islands, being lower at the centre than in the north-east and west. Its height is about 750 feet and it is covered with trees except at those places on the hill sides where large outcrops of rock occur. On such spots what vegetation exists is of a sparse and stunted type.

For some distance from the shore a reef filled up the bay where, indeed, the conditions are most favourable for the growth of corals. As one rowed over the pellucid green water, looking down they were to be seen in indescribable variety—great heads formed like massive boulders and tiny sprays no less delicate than a piece of moss. Corals of all shapes and shade were there—pink, grey, yellow, brown, blue, green, red, while among the crevices and branches swam fish as gorgeous as their surroundings—little fellows half an inch in length, blue, red, and yellow and others of larger size whose brilliancy of colouring passed almost unremarked by comparison with the grotesqueness of their forms.

In the centre of the bay and connected at low tide with the shore stood a rocky islet frequented by numbers of the white tern (*Sterna bergii*) with rose-tinted breasts from which the flush fades immediately after death.



From several small streams within the jungle good water may be obtained; they are, however, too weak to force a passage to the sea and must be looked for above the beach.

At Pengiki we spent two days but found the island, although of fair size—three miles long and nearly two wide—very scantily furnished with animal life. Pigeons both Nicobar and *bicolor*, were fairly numerous as were the glossy starling and blue-and-white kingfisher, but no other birds were seen with the exception of an eagle and a single grey wagtail. A python and viper were obtained and a glimpse was caught of a small animal in a tree that may have been a musang. Macaque monkeys (*cynomolgus*?) were, however, common and a new species of squirrel (*Sciurus mimivulus*), a dwarf form of *S. prevostii*, was obtained.

The jungle was of fair height but possessed scarcely any undergrowth and all day long in the cool green light, swarms of bats (*Hipposideros barbensis*, sp. nov.) flitted about circling in and out amongst the trees.

When we left—in darkness at three o'clock one morning—the anchor proved to be jammed fast in the coral. It was only by setting full sail, after all other means were exhausted, that we were enabled to break it free.

#### The Tambelans.

The Tambelan group, about sixty miles to the northward, was the next place of call. With a wind abeam and a squall to help we arrived in twelve hours and anchored between two small islands, Selendang and Gilla. Exploring the former in the afternoon we got two or three birds. It possesses a curious cone-shaped hill about 700 feet high; Gilla is much lower. The scenery in the little strait between the two was very lovely—jungle and coco palms, granite boulders and yellow sand, bright blue sea and waves of white surf at the far end of the passage.

Early next morning (Aug. 4th) we boated over to Great Tambelan and met many canoes going off to the outlying islands. The sea was running very high a few hours later and the sail back again was somewhat exciting.

The Tambelan group consists of the three principal islands of Tambelan, Bunoa and Wai, with a number of smaller islands



massed fairly close together over thirteen or fourteen miles of sea. Only Great Tambelan is inhabited but on the others the people have numerous *ladangs* and *pondoks*. The population consists of Malays, 500-600 in number.

### Bunoa.

As Gilla and Selendang were too small to be productive, after one more visit we transferred operations to Pulo Bunoa close by. The islands form roughly two parallel lines lying N. E. and S. W. Bunoa is the largest of the south-western group. It is about four miles long and two and a half wide, on the whole gently rounded in contour with gradual forested hills rising to a height of 900 feet. On the northern side is a bay that offers good anchorage in the S. W. monsoon.

We tramped through the island for three days after birds and beasts. A form of "Krah" was common and from those collected here, and later on in the Anambas, a new species *Macacus punilus* has been described which differs from *M. cynomolgus* in being paler and much smaller. The little pied hornbill (*Anthracoceros convexus*) was numerous: their chattering resounded through the jungle like the yelping of puppies. Once in stalking them I lost my bearings and at last crossing a slight trail followed it up on the wrong hand and passing by a little stream that suddenly disappeared subterraneously, came on a ladder path and log-slide running down a steep hill side to a strange beach that in the end proved to be on the further side of the island. Thinking it easier to return to the boat by following the coast than by retracing my steps, I let myself in for a five or six miles scramble in mangroves and mud, over soft sand and rocks, across coral reefs and through water breast high before I found the schooner again.

Arenga palms are numerous in the jungle and are worked for sugar by the natives. The trees are tapped near the top and bamboo receptacles are fastened beneath the incisions to receive the sap. Here and there we came across the boiling-down furnaces that consisted of large flat iron pans raised above the fire on clay walls. Before concentration the sap is carefully strained through a bunch of fibre to remove impurities and is then boiled down either to a treacly consistency or to a still

greater density when it is poured into bamboo moulds and crystallizes into a dark brown sugar.

A coconut shell of the warm sweet liquid in its early stage is most refreshing during the course of a hard tramp and was always offered when we passed a concentrator at work.

### Great Tambelan.

On the 8th of August we made sail at day-break and with a native pilot crossed to Tambelan Island and anchored in the harbour three hours later.

This, the largest island of the group, is nearly  $4\frac{1}{2}$  miles long and roughly triangular. Near its north-east coast are several hills, the highest of which—Tambelan Peak—rises to an elevation of 1300 feet while a short distance to the eastward of it is Thumb Peak, a remarkable pointed hill 950 feet high. The island is nearly divided into two parts by an inlet of the sea which runs in a north-easterly direction into its western side. This creek is nearly a mile in width but is fringed and choked with reefs. The remains of a breakwater built of coral cross it about a mile and a half from the entrance. A stockade once ran behind this and a fort stood on the shore, all being constructed to defend the village which lies higher up, from the attacks of Illanum pirates who occasionally visited these islands years ago.

We threaded our way amongst the coral and anchored in a clear patch of water near a couple of small native schooners, well protected by reefs from any south-westerly swell and in sight of the Kampong. The conditions permitted some delightful baths in perfect safety for the Malays said crocodiles and sharks never ventured into the neighbourhood. The pilot refused money payment for his services but gladly received drugs for an ailing relative, and later when the people took to bringing us specimens they always preferred medicine of sorts by way of recompense.

We first landed on the southern side of the harbour and climbed a small hill of granite and laterite. It was covered with scanty scrub and absolutely devoid of life. A reward however lay in the view. Below the slope of the hill stretched the still green waters of the harbour, purple-patched with coral; on the further side sand, jungle and palms, while the Kampong—a



long line of brown houses on land and water, following the course of the shore—lay higher up the creek. Beyond rose the jungle and hills, with the quaintly-shaped Thumb Peak—abrupt and solitary—standing out prominently against the sky.

A white man is a bit of a curiosity in these parts and for the rest of the day it was but seldom that twenty or thirty heads were not poked through the skylight windows watching our goings-on.

Next morning we went up to the village to visit the Dato and show him a letter written by the Sultan of Linga who, under the Dutch, rules all these islands.

A roofed-in platform with benches and flight of steps forms a pleasant landing and lounging place before the village, near which was anchored a stranger vessel—a little prau about the size of our whaleboat packed with a large and unprepossessing family of Orang Laut.

From the *jambatan* we were led to the Dato. The village consists of about 250 houses threaded by tidy paths and supplied with water by numerous bamboo *penchurans* leading from the higher ground in the rear. No women were visible but from the houses came the clacking of numerous looms: good strong sarongs are made here, dyed, however, with aniline colours purchased in Singapore: indeed though Dutch possessions, the Straits dollar was the only coin current in all the islands of our cruise. Passing the mosque, a pucca triple-roofed structure of wood and stone, and a number of old cannon that lay scattered about, we reached the Dato's, a well-built shingle-roofed house, with a long enclosed verandah running its length, in which we were welcomed.

The Dato was an amiable old gentleman, treated by his people as he treated the Sultan's letter—with great respect. Chairs were arranged at one end of the room and when we took our seats the lower end was crowded by the populace while women stared through the latticed windows of the inner rooms. An official, bent double with respect, read the Sultan's *hukum*, which explained our object and recommended all assistance, and then the Dato and the audience descanted on the local fauna, its paucity in those islands and how much better off other places—Pahang, for instance—were in this res-



pect. On leaving, the hand-camera was brought into play for the first time and all the juveniles among our escort fled screaming.

The little community appeared to be very well off and was one of the few places remarkable for the absence of the ubiquitous Chinaman, a solitary trader of which race seemed to have been recently starved out. On the shores of the harbour twenty to thirty schooners of local construction were drawn up: these seemed to be owned by the villagers in general, and with them all the trade was done and all the produce shipped from time to time to Singapore direct.

Ship-building bulked large as an industry of the village and we saw half a dozen or more hulls from thirty to forty feet in length, in all stages of construction. These vessels are built of locally grown chengai and merbau and are fastened with tree-nails throughout. Each seemed to be the work of about a couple of men in partnership and takes two years or so to complete. A few frames are first set up and completely planked and afterwards the other ribs are fitted in until sufficient strength is attained. It was said that a 35 foot craft (10-tonner) could be purchased all complete for \$350, and although perhaps their lines were capable of some improvement they were fine roomy little boats strongly built of thoroughly sound material. The local canoe, however, was a thing of beauty: strongly built of two prettily contrasting white and brown hardwoods without a single nail, with upward-projecting stem and stern and gaudily painted bird's-head bracket on the bow to support the lowered sail and mast, it was as workmanlike as it was handsome. The sail was a square cotton lug slung by the middle of the yard and set with the forward end of the boom snubbed down to the lee-bow well forward.

Having sent the Dato a photograph of the "Terrapin" we received a call one evening from him and his understudy to acknowledge the picture and to obtain a little medicine. Amongst other things he told us how in his boyhood the village suffered the last attack from pirates and how all the inhabitants were driven out by the raiders to a hill at the back of the island where they built a fort of refuge. In those days there were only about a hundred people on the group.

But little was to be obtained by way of supplies from the village. Coconuts and copra were plentiful, eggs and fowls scarce: we could get plenty of bananas and one evening bought from a canoe homeward bound from fishing, three splendid parrot fish (*Scarus sp.*), weighing together between sixty and seventy pounds for a dollar!

The jungle was intersected with paths leading to the arenga palms and trying-down sheds and by shooting along them and in the cocopalms we obtained a new squirrel (*Sciurus abbotii*), a pale form of *S. notatus*.

After investigating the birds and mammals on several occasions with good results, we devoted a morning to butterflies getting about a dozen species round the village and along the forest paths. The fauna of the island was neither numerous nor diversified and on the morning of the 15th we moved the schooner over to Pulo Wai, anchoring off its N. E. coast.

#### Pulo Wai.

This island is the most north-westerly of the group. It is about two miles long and rises in several peaked hills attaining near the eastern end a height of 1000 feet. Being farther from the Kampong than the others it is least visited but plantations of coconuts and bananas, plantains, yams and sweet potatoes are common on its hillsides, a good deal of which are cleared.

It provided us with a handsome squirrel (*Sciurus mimellus* sp. nov.) with black, chestnut and white pelage—a dwarf form of the well-known *S. prevostii*.

A walk across the island proved very bad travelling but from the hills a distant view was obtained of Gap Rock about twelve miles to the northeast. This remarkable islet consists solely of two huge boulders—the larger of which is 124 feet above the water—lying on a flat platform of rock utterly devoid of soil or vegetation.

This was the last of the Tambelans visited and I will therefore conclude this notice of them with a list of the principal collections.

#### Mammals.

1. *Macacus pumilus*, sp. nov.

2. *Pteropus lepidus*, sp. nov.
3. *Megaderma spasma*.
4. *Tupaia bunoe*, sp. nov.
5. *Tragulus* sp. (said to occur).
6. *Sciurus memellus*, sp. nov.
7. *Sciurus abbottii*, sp. nov.
8. *Mus tambelanicus*, sp. nov.

#### Birds.\*

1. *Gracula javanica*.
2. *Eulabes javanicus*.
3. *Calornis chalybeus*.
4. *Hypothymis azurea*.
5. *Hirundo javanica*.
6. *Motacilla melanope*.
7. *Halcyon chloris*.
8. *Anthrococeros convexus*.
9. *Cypselus* sp.
10. *Osmotreron bicincta*.
11. *Carpophaga aenea*.
12. *Myristicivora bicolor*.
13. *Chalcophaps indica*.
14. *Calænas nicobarica*.
15. *Turtur tigrinus*.
16. *Esacus magnirostris*.
17. *Totanus hypoleucus*.
18. *T. calidris*.
19. *Lepterochus sacer*.

#### The Rocky Islets.

Eleven miles in seven hours is not good sailing but it was afternoon when we landed on the Rocky Islets—the Pulo Mandariki of the Malays. They consist of two small barren islets and a rock. The sea is steep to all round and the schooner lay on and off with a kedge anchor hanging down while we care-

\* It is interesting to note that no birds smaller than the fly catcher have effected a lodgement on the Tambelan group.

Jour. Straits Branch



fully humoured the swell and jumped ashore on the larger of the two, which is northernmost. The central islet which is next in size is much more broken and is a wild confusion of little peaks and precipices.

Climbing to the top of our islet (134 feet) we found that quartz predominated in its formation, which presented an exceedingly rough surface where grew here and there only the scantiest tufts of coarse grass. Two or three dead bushes crowned the summit and everywhere the ground was fouled with the guano of sea-birds that use the rocks as a breeding place.

The air was full of squalling, squawking, croaking gulls and among the crevices of the island's surface we discovered numbers of nestlings, and a few eggs all addled. The birds were of two kinds:—a black noddy with grey head (*Anous stolidus*) and a tern (*Sterna* sp.) with black head and wings, grey back and grey-white breast. The remaining fauna included fishing-eagles, crabs and a solitary bee. The mid-day sun, shining down on the bare rocks, made the heat, combined with the scent of the birds, overpowering and we soon left, having knocked over sufficient specimens for the collection in a very few minutes.

### Saddle Island.

Instead of proceeding straight to the Anambas we stopped for a night and morning at Pulo Kayu Ara, a little island about half a mile long covered with jungle and thus contrasting strongly with the place we had just left. It is nearly 400 feet high and is formed of two round hills having a dip between that make together a contour from which it has gained its English name.

The feet of the hills were fringed with black rocks but between lay a delightful little beach inhabited by a pair of white-collared kingfishers, on which we landed and found the laying places and tracks of turtles: the sea, of wonderful limpidity invited to a bath. The only birds seen in the forest were fruit pigeons and glossy starlings,—the only terrestrial mammal a squirrel (*Sciurus klossii*), a small blue-bellied member of the *notatus* group. The trees bore considerable quantities of fruit,

particularly noticeable being wild nutmegs and mangosteens, the last of which seemed to form the principal food of the squirrels. To complete the tale of our acquisitions were the small fruit bat (*P. lepidus*) previously taken in the Tambelans, some geckoes and two or three tree-snails.

Waiting until the tide turned in one favour we left in the afternoon, bound for the Anambas, a hundred miles away. Until sunset next evening the wind held light but then with a series of squalls coming up from astern we made from six to seven knots an hour, the breeze constantly shifting a point and back and compelling us to make continual gybes. Late at night we entered the channel between Pulos Peling and Riabu and dropped anchor close to the former.

#### The Anambas Islands.

On the morning of the 17th it was evident that we were fortunate to have anchored when we did; for dead ahead, just a hundred yards away, an abrupt bit of coral reef on which the schooner could have piled up very awkwardly, lay five or six feet beneath the surface.

Pulo Peling, which we worked for a day, is only a small island without inhabitants where clearings were just being commenced; we saw no mammals except many monkeys, but got a little blue brown flycatcher (*Cyornis tickelli*) with reddish breast for the first time on the cruise.

Riabu, which we next prospected, is much larger, being high and about six miles in length. It possesses a very good bay affording excellent harbourage in the S. W. monsoon as it is only open between N. and W. As we rowed along the shore we shot a large "baiawak" from the boat and then landing found the country very rocky and densely covered with forest. Our best catch was a squirrel—the only one seen—a pale and rather small form of *Sciurus notatus*. It has been named *anambensis* and occurs on most of the islands of the Anambas group but the Riabu specimen differs from the others in having the audital bullæ markedly smaller.

At daylight on the 19th there was scarcely a breath of wind and after getting up anchor we nearly drifted on to the reef



through which we had so nearly come to grief before, but missing it with the rudder by a bare yard we soon afterwards got a fair breeze and, running past eight or ten small islands, made Terempa, which is the chief kampong of the Anambas, by 2 p. m.

The Anambas Islands are situated in lat. 3 N., long. 106 E. and extend over an area 65 geographical miles long N. and S. and 55 miles wide. They practically form two groups of which, though the western includes the larger island, the other is greater in area. All the islands are hilly, covered with forest, and afford numerous bays and channels where safe anchorage may be found in spite of the coral reefs that occur everywhere. Although the population is mainly confined to the larger islands, Malays are thinly scattered over most of the rest and there are Chinese Settlements in either group.

#### Siantan.

Terempa lies in a little bay on the north coast of Siantan which is the largest of the easterly Anambas, having probably an area of about 20,000 acres: it is densely covered with heavy forest, is very hilly and rises 1855 feet.

The kampong is a thriving little place with a Chinese cemetery, twenty or thirty Chinese shops with galvanised iron roofs and a good proportion of its population Chinese. A small steamer the "Banka" calls once or twice a month and yet it is against these islands that the dear old China Seas Directory (3rd edition) still contains the antediluvian warning that "it is dangerous to land without due precaution, for the Malays who reside on them may probably massacre or make slaves of strangers if they perceive a convenient opportunity." *A propos* of this sort of thing I remember once asking a Danish sailor whom I met on the other side of Sumatra whether he used the English Admiralty publications for these seas, but the skipper of that very old-fashioned little barque the "Hans of Fano" shook his head and replied that he always consulted certain continental sailing directions as our own were far too obsolete and scrappy.

Terempa is built along the head of the bay and has a small saltwater creek opening out behind it. As one faces it from seaward an orderly street of Chinese stores with the opium-farmer's place and a Chinese school lie to the left with a few



Malay houses at the far end. In the centre is the Dato's house—a wooden building in civilised style; a large house belonging to the Sultan of Linga and then stretching away on the right the houses of the bulk of the Malays: behind the flat on which the village is built, hills covered with jungle and coco palms rise steeply in a half circle, the whole having a very pretty effect which, however, is quite matched by the view presented from behind the town from whence looking down the long slope of the hill one sees the roofs of the houses, the semi-circular bay with boats lying at anchor, and then across a strip of water the forested hills that rise opposite.

A mile and more to the north of Siantan lie the two fairly large islands of Mobur and Mata with a channel about a mile wide between them. Between Mata and Siantan runs a very pretty strait which although obstructed by many islets and shoals at its eastern end, has deep water in the west where Terempa Bay is situated. Not only in the latter itself can a vessel lie in deep water sheltered from all winds, but beyond the point forming its eastern shore is another bay where a ship can anchor in 10-15 fathoms near the beach, entirely land-locked. Better harbours on a small scale could not be wished for and wood and water are close at hand.

The Dato of Terempa was in an advanced state of consumption and in spite of possessing a new and rather showy house lived in a second more modest dwelling. He showed us specimens of black iron ore of which large quantities were said to occur in the hill behind the village.

We could get various supplies here, thanks to the presence of the Chinese; extremely good coconuts\* were plentiful and water was at hand in a large pool on the beach. The only practicable shooting ground near at hand was mostly covered with small scrubby jungle in which we shot squirrels and a number of fairly common birds and trapped plenty of rats. I met with

\*This group of islands would appear to be a convenient place for obtaining seed nuts to form a coco-palm plantation. The Malays always maintain, and apparently with some show of reason, that island nuts are far superior to those of the mainland. Some of the Anambas nuts are very good, in fact, almost equal those of the Nicobars: though small, they are extremely sweet and the deposit of flesh is very thick.

bees in this place and was put out of action for a day or two in consequence, for while chopping at a liana that crossed the path I disturbed a swarm that were clinging to it a little to one side. They were about me in an instant and though by retreating to the sea at an unaccustomed pace I shook off the majority, nevertheless those that had secured a hold made things sufficiently painful and my right hand and arm were so swollen that for a time I could do nothing with them.

The canoes of these islands were very similar to those of the Tambelans; they differed principally in having a much higher stem and stern and were shallower, lighter and narrower, with much less beam forward than aft, and they were propelled with a double-bladed paddle. The sampan used by the Chinese was of a very bulky model with an elongated and upward pointing prow: it sailed under a battened dipping-lug.

Having been told by a son of the Sultan of Linga who was staying in the village, of a large waterfall on the east side of Siantan we set out at sunrise one morning to visit it in the whale-boat. The wind was dead ahead and we had to pull all the way—about eight miles—but the result was well worth the labour. The way lay right up the channel between Siantan and Mata, then a short distance down the east coast and finally an approach by a passage through mangroves that enabled us to bring the boat to the very foot of the falls themselves. Bordered by jungle these fell in a series of cascades down a bare strip of rock about 400 feet high. There were twelve or thirteen falls in all and their chief charm lay in their variety—broad ribbons of foamy water slithering over smooth faces of rock, long threads dropping uninterruptedly from a height, and series of little cascades tumbling down flights of stone steps, while here and there in between were delightful cool-looking pools, seeming so inviting after our hot row that we felt compelled to have a swim, following the example of the fish that had somehow found their way there.

The raja had agreed to accompany us but was late in setting out, though he arrived before we left and we both started together to sail back, as this time the wind was fair. We were not long in company however, for the other boat—curiously, built in Singapore in the same yard as our own—a little larger



and with more sail soon left us behind and we reached home badly beaten.

The days between August 24th and September 5th were passed in visiting the northern island but on the latter date we sailed round Siantan and the islands extending from its south-east extremity since the channel on the north was untraversable, drifting on a reef on the way in a calm but easily getting off by the use of a kedge anchor, and in the evening putting in at Telok Ayer Bini, a bay on the south coast partially protected by an island at the entrance.

The shores rose very steeply and were uncleared except in one or two places where the people of a house there had made gardens and planted hill rice. It was hard work climbing about the slopes which were rendered excessively slippery by rain that continued incessantly throughout our stay. A stream with two arms ran into the head of the bay and up one we rowed until stopped by a small cascade, while the other was merely an almost dry bed of granite boulders.

The wretched weather made specimens scarce and after three or four unprofitable days we sailed on our final visit to Terempa. Four more days were spent here working fresh ground and making fairly good collections of mammals and birds and then we left for Pulo Telaga to the westward.

#### Mobur.

The 24th of August was the first day of our stay at Mobur Island, about 5 miles to the northward of Siantan. Like all the Anambas it is hilly with a broken surface covered with forest, and on the south it is cut into by a narrow gulf with an islet at the entrance; a notable feature of the western group is the number of these inlets and narrow channels that occur. Good as the harbour was with depths of 12 or more fathoms we passed up the strait separating Mobur from the larger Mata and anchored in a big bay on the north side, landlocked for more than three quarters of its perimeter. Four or five miles seawards we could perceive the Tokong Belauer, a remarkable white rock bearing at a distance a most extraordinary resemblance to a modern battleship. We found coco palm and banana plantations on the island but very few inhabitants, for the people



are mainly confined to Terempa and only isolated settlers occur elsewhere.

One afternoon we rowed to a little island in the bay called Langor. It was only about 100 feet high and consisted of sand and rocks covered with scrub and a few cocopalms. Everywhere the ground was strewn with pigeon's feathers and by waiting we found that large flocks of the orange-breasted pigeon (*Osmotreron bicincta*) came off at sunset from surrounding islands and used this spot as a roosting place in company with smaller numbers of glossy starlings and nutmeg pigeons. The whirring noise made by the wings of the flocks as they flew round and round the island, disturbed by our shots, was very great and continued until we departed with bags stuffed with birds that appeared later in a most delicious stew.

#### Kelong, Manguan and Tobing.

We next sailed round the north end of Mata and anchored between it and Pulo Kelong, a narrow island about 5 miles long in a N. and S. direction and less than a mile wide: the ground sloped upward to a ridge 600-700 feet high and every where the soil and jungle were very poor. Most of the channel dividing Kelong from Mata is filled with sand banks and coral, dry at low tide. Fishing-stakes had been set up here and there by the Malays but to us the place proved a good ground for the common shore birds of this region and we also collected a number of beautiful starfish while our crew hunted for trepang and chopped tridacnas out of the coral. Other animal life was scarce and we soon moved southward to Pulo Manguan, a small island shaped like a dumb-bell, flat and swampy in the centre; but doing no better there, anchored the schooner off Tobing, an islet near the eastern entrance of the Siantan channel, and from thence again visited the waterfall, and next day, previous to sailing for the south of Siantan, rowed to Terempa and back for our mails.

#### Telaga.

The second stay at Terempa concluding on the 13th of September we made for Jimaja, the chief of the westerly Anambas, first however after a few hours sail stopping about mid-way, at a group of small islands of which Telaga is chief. This is a

narrow hog-backed island about 5 miles long N. and S. with a picturesque peak rising 1740 feet near its northern end. We spent a couple of days here and on the neighbouring island of Midai finding the forest fairly open except for patches of rattans and a prickly holly-like shrub. Only one village of three or four houses was seen. The coconuts seemed much troubled with squirrels and the people made use of an ingenious trap set on a long bamboo leading from the jungle to the palm trunks.

### Jimaja.

On the 16th we made sail at daybreak and rounded the southern end of Telaga. The wind was ahead but we did most of the seventeen miles, which is the distance between that island and Jimaja, in one board and then working short tacks got close to the entrance of Kwala Maras Bay on the east coast by night-fall. After that the wind fell light and the tide carried us away to the northward so we let go the anchor in 10 fathoms about three-quarters of a mile from shore and next morning after a couple of hours beating got into the bay. A line of rocks extends above water from the north shore and beyond them on the other side is a coral reef: we found a good berth between the two, well protected from all but easterly winds.

Jimaja is the largest of the Anambas with an area of perhaps 30,000 acres. It is of an irregular Y-shape about 14 miles N. and S. and 9 miles wide. The contour is very uneven and there are many peaks between 700 feet and 1530 feet, the greatest elevation attained in Gunong Tujoh. The irregularity of the coast line has resulted in many bays, the largest being in the north and in the south-east, but Telok Kwala in the centre of the east coast, although smaller than these is the most important as having the greatest population and being the port of call of the steamer. Its shores rise steeply to heights of 1000 feet in places free from forest, but beyond the head where a river embouches is some flat swampy land overgrown with mangroves.

The kampong lies on the north shore and consists of thirty to forty houses, a small mosque, the Dato's house and the buildings of the opium-farmer—shop, godowns, etc., surrounded by a stockade 8-10 feet high closely built of small saplings from whence ran a small jetty.



The Dato's was a well built house, the largest in the place, with a flag staff in front. We were given the rarely used chairs brought out, as always, from somewhere in the roof and seated on these awaited the Dato who was making himself presentable. He was a rather big man with an Irish countenance and wore a "baju tangan kanching" having a ridiculous resemblance to the obsolete night shirt, and a purple smoking cap whose large black tassel persisted in lying in his right eye. There were only a few people in the audience and after the Dato had stumbled through the Sultan's letter and the scholar of the party recorded our visit in the brown-paper-covered archives of Jimaja, conversation took a zoological turn and we were told also of two deep lakes with waterfalls that existed up in the hills of the interior, the description being such as to give one an idea of something impressive.

We collected first on the south side of the bay with no great result; the ground was very steep and difficult and the forest small having evidently been cleared at some former period. However, a new species of rat—*Mus flaviventer*—was obtained and we came across parrots (*Paleornis longicauda*) for the first time. Every day flocks crossed the bay and passed near the summit of some steep rocks where I several times lay hid in the vain hope that they might stop. It was interesting to note how the parrots' call as they flew by always drew a loud response from all the small birds roosting in the bushes.

On the other side of the bay we found a good path running for some distance through a former mangrove swamp now planted with coconuts, and afterwards up hill amongst most beautiful jungle. It crossed two small streams of perfectly clear water one running among granite boulders and the other, about a foot deep, in a smooth sandy bed. Toward midday nothing was more pleasant after five or six hours' tramping in the forest than to find a round stone for pillow and recline full-length in the flowing water of the latter until thoroughly refreshed by its coolness. But beside providing pleasure of this sort we also obtained good specimens along its banks.

Our third collecting ground was along the river falling into the head of the bay. At first this was bordered by broad growths of tall mangrove, but after a mile or so where the current



began to make itself felt these ceased and the vegetation became more varied with interminglings of nipahs, palms, and fruits trees. Onward from here the stream—the Sungei Maras—runs up the centre of a winding valley with a narrow strip of flat land at the bottom and steeply rising sides. At the head of navigation—and in fact when the tide was low we had to wade with our dinghy for some distance, though all the way the stream was broad enough to use oars—was a village of eight or ten houses and a mosque, one or two of the former being of large size and well built of panels of carved wood, though now old.

In the valley we got specimens of the parrots at last as they fed in the fruit trees bordering the river, and also a lovely little two-toed kingfisher (*Ceyx rufidorsa*) with coral red beak and feet, yellow breast and brick red head and back that were tinged with a beautiful glossy lilac: the best of the bag however were specimens of a big squirrel (*Ratufa anamba* sp. nov.) black above with tawny yellow cheeks and underparts. It weighs about 3½ lbs with a total length of 33 inches of which head and body are 15 inches: thus, contrary to the general rule, it is an insular race characterised by increase of size. In these China Sea islands it is found that when a species of mammal occurs in a solitary island, however small that may be in area, it is confined to that island alone, but when a species is present on an island of a group it will generally be obtained throughout the group. In the case, however, of the *Ratufa* of Jimaja this is not so; it was neither reported nor did we find it ourselves on any other of the Anambas.

On the 25th we set out early in the morning with the intention of seeing what truth there was in the report given us of the "telaga" in the centre of the island. The first stage was to the village up river—Kampong Ayer Maras—where the Penghulu provided us with a guide. While waiting for the latter we inspected a small waterfall about 20 feet high at the back of the village. It was not much to look at however as there was very little water in it at the time. We were then told that there were two series of lakes, so voted for the larger set and when the guide arrived started him off accordingly.

The path traversed sago swamps for the greater part of the way and was very muddy; then passed through a small kampong surrounded by fruit trees and soon after that through a stream

where a sago-making apparatus was erected, next came thickets of dense scrub followed by another sago plant where a number of men were at work. Now the track degenerated into a muddy ditch knee deep for the most part and after crossing several brooks we came to the bank of a small river with a clean sandy bed. When we had waded upward for some distance the bed became rocky and we then soon reached the "lakes." These were disappointing being merely large rocky basins in the river bed about 60 feet long and 30 feet wide and, as we found by diving, 17 or 18 feet deep. They were connected by a fall and there was a second above the upper pool both some 30 or 40 feet high. The water was perfectly clear and the whole very pretty, rock and water being shaded and hedged in by dense jungle, nevertheless the actual state of affairs was not quite the phenomenon it had been painted by our informants.

We found the steamer in the bay when we got back and her *serang* came off with a message from the commander that we had stolen his anchorage! As however we had been there some time and the other was still under steam we returned word that we felt no inclination to move; thereupon the steamer's master obstinately took up a berth a few yard off until he swung with the tide when, our main-boom end doing considerable damage to his bridge dodger, he was persuaded to seek a more convenient anchorage.

Both in the Tambelans and in Siantan we had made efforts to get one of the canoes of the place without avail, people did not want to sell or would not be ready in time. Here as a last chance one of the built up kind was to be had for \$40, though graceful dugouts on exactly the same lines were just half that while rough models could be purchased for as little as \$3. One canoe, a good example of the type—was brought alongside with the sail lightly rolled up and bound round and round with every conceivable cord; undoing this tangle of course showed the cotton to be mildewed and full of holes and the chagrined vendor was sent off for another. There was further trouble in concluding the purchase as the islanders would not accept either Dutch coin or Singapore notes and we had run completely out of Straits money. Happily a couple of Tringanu men visiting the island in a small prau were willing to change our notes on condition that

we invested in sarongs from their trade-stock. Having with the assistance of these men—who contrasted greatly with the islanders in many ways—brought the canoe business to an end we left Jimaja on September 28th and sailing round the south of the island reached Singapore on October 8th via Tiunan where we spent a few days.

The principal result of our visit to the Anambas lay in the first record of the animals and birds found in them, an outcome of our investigations being also the description of many new insular species. So far as is represented by the collections the mammal fauna of these and of the other islands visited consists of local forms of the widely distributed and characteristic Malayan types. Each island and group of islands has its representative of the common genera and species; but in scarcely an instance is an insular race identical with that occurring on another island, unless of the same group, or on the mainland.

#### Mammals of the Anambas.

*Macacus pumilus*.  
*Nyctcebus tardigradis*.  
*Emballonura anambensis*, sp. nov.  
*Rhinolophus minutus*, sp. nov.  
*R. rouxii*?  
*Tupaia chrysomella*, sp. nov.  
*Paradoxurus*, sp. (reported).  
*Tragulus*, 2 sp. (reported).  
*Sciurus anambensis*, sp. nov.  
*S. tenuis*.  
*Funambulus castaneus*, sp. nov.  
*Ratufa anambæ*, sp. nov.  
*Mus siantanicus*, sp. nov.  
*M. strepitans*, sp. nov.  
*M. anambæ*, sp. nov.  
*M. flaviventer*, sp. nov.

#### Birds.

The birds obtained were all common peninsular forms and are enumerated below.

*Malacopteron magnirostra*.



Anuropsis sp.  
Mixornis gularis.  
Ægithina viridissima.  
Pycnonotus simplex.  
Iole sp.  
Dissemurus platurus.  
Orthotomus atrigularis.  
Lanius cristatus.  
Eulabes javanensis.  
Calornis chalybeus.  
Hypothymis azurea.  
Cyornis tickelli.  
Cittocincla macrura.  
Munia semistrata.  
Hirundo gutturalis.  
H. javanica.  
Motacilla melanope.  
Anthothreptes rhodolæma.  
Æthopyga siparaja.  
Æ. hasselti.  
Dicæum trigonostigma.  
Halcyon bengalensis.  
H. chloris.  
Ceyx rufidorsa.  
Cypselus subfurcatus.  
Collocalia fransica.  
Macropteryx longipennis.  
M. comata.  
Rhamphococcyx erythrognaethus.  
Graculus sumatrensis.  
Palæornis longicauda.  
Loriculus galgulus.  
Spizaetus or Spilornis sp. (observed).  
Haliæetus leucogaster.  
Osmotreron bicincta.  
Carphophaga ænea.  
Myristicivora bicolor.  
Calænas nicobarica.  
Chalcophaps indica.

*Charadrius fulvus.*  
*Ægialitis* sp.  
*Esacus magnirostris.*  
*Totanus calidris.*  
*T. hypoleucus.*  
*Strepsilas interpres.*  
*Tringa hypoleuca.*  
*Fregata aquila* (observed).  
*Anous stolidus.*  
*Sterna bergii.*  
*S. media.*  
*Leptorodius sacer.*  
*Butoroides javanicus.*

#### Literature.

China Sea Directory Vol II.

Mammals collected by Dr. W. L. Abbott in the region of the Indragiri River, Sumatra, by Gerrit S. Miller, Jr., Proceedings of The Academy of Natural Sciences, Philadelphia, March, 1902.

Mammals collected by Dr. W. L. Abbott on Islands in the South China Sea, by Gerrit S. Miller, Jr., Proceedings of the Washington Academy of Sciences, August, 1900.

# A List of the Butterflies of Borneo with Descriptions of New Species.

BY R. SHELFORD, M.A., F.L.S.,  
(Curator of the Sarawak Museum).

## Part I.

(DANAINÆ TO AMATHUSINÆ).

Only three lists of the butterflies of Borneo that can pretend to any completeness have been published hitherto. One by Messrs. Distant and Pryer appeared in the "Annals and Magazine of Natural History" 1887; another by Messrs. Pryer and Cator in the "British North Borneo Herald" of 1894—a list remarkable for the number of *nomina nuda* contained in it, and a third by Mr. E. Bartlett in the "Sarawak Gazette" of 1896, reprinted in the "Zoological Note-Book of Sarawak" by the same author. All three lists are now out of date and many new species have been described even since the compilation of the last one; for example Mr. Bartlett records only 139 *Lycænidae*, whereas the number of species of this family now known from Borneo just exceeds 300. Consequently I feel justified in adding to the literature on Bornean *Lepidoptera*, especially as I do not expect that many more discoveries of new species will be made in the future in an island that has been so well explored faunistically as Borneo; in short, this list has some claim to completeness and finality.

A few new species are described for the first time, and as considerable trouble has been taken in consulting all available literature on Oriental butterflies and as most of the specimens have been referred to Dr. A. G. Butler and Mr. F. A. Heron, of the British Museum, and by these well-known authorities de-



clared to be in every probability undescribed, I trust that my new species are really "good species."

The question of nomenclature has been rather a difficult one; the systematist has to steer between the Scylla of "lumping" and the Charybdis of "splitting," for to regard all geographical varieties (topomorphs) of a wide-spread species as identical is unscientific, to regard each variety, on the other hand, as a separate species is almost equally unscientific and, further, tends to obscure the problems of geographical distribution. There is a third alternative and that is, to adopt the cumbersome trinomial system, distinguishing well-marked topomorphs of a wide-spread species as sub-species, giving them separate names\* and noting the distribution of the species as a whole. Such is the method adopted in this list; a species is recorded, if the typical form occurs in Borneo that fact is noted and the distribution of the typical form and the sub-species (component parts of a species) occurring outside Borneo are also noted; if the typical form does not occur in Borneo, the name of the sub-species that does occur follows the name of the species and the distribution of the species as a whole is recorded.

Unfortunately some of the families of butterflies have not been studied so thoroughly as others, whilst some families again have been studied from a different point of view to others: for example, Messrs. Elwes and Edwards in their "Revision of the Oriental Hesperiidæ" (Trans. Zool. Soc. Vol. XIV part IV, 1897), do not allow a single sub-species, whereas Hon. W. Rothschild and Dr. K. Jordan in their memoirs on the Papilioninæ and on the Nymphaline genera, *Eulepis*, *Charaxes*, etc., regard every topomorph of a wide-spread species as a sub-species and this to my mind appears to be the more scientific method. Much work along these lines still remains to be done, but it can be done only by one who has frequent access to extensive collections and well-stocked libraries, therefore though this list lays claim to some completeness so far as an *enumeration* of the forms of butterflies

\* For an able exposition and defence of the trinomial system of nomenclature see *Novitates Zoologicae* Vol. IX. Supplement, 1903, pp. xxvi et seq.

occurring in Borneo goes, it does not claim in every instance to discuss the relationships between these Bornean forms and close allies inhabiting other areas.

At the end of this list will be given a table showing at a glance the geographical distribution of the various species. A few field notes are put in square brackets after the records of distribution of the respective species.

I am indebted to Mr. F. Moore and to Herr H. Fruhstorfer for their kind help; the late Mr. L. de Nicéville was a most valued correspondent whose sound advice and generous aid I sadly miss now.

#### LEPIDOPTERA RHOPALOCERA.

##### Fam. NYMPHALIDÆ.

##### Sub-fam. DANAINÆ.

##### Genus *Hestia*.

##### 1. *Hestia lynceus*, Drury.

*H. lynceus*, Drury. Illustr. Exot. Ent. ii. pl. 7 f. 1 (1773).

*H. reinwardti*, Moore, P. Z. S. 1883, p. 218.

Sub-sp. *H. lynceus druryi*, Moore, with a melanic form *fumata*, Fruhst.

Fruhstorfer (Berl. ent. Zeitschr. Bd. xlii. p. 314. 1897) divides *H. lynceus* into five sub-species occurring in Malacca, Penang, Singapore, Nias, Sumatra, Mentawai Is., Natuna Is., Borneo and Java. The species is a common one and the Bornean sub-species with its melanic form is distributed throughout the island; *H. lyn. druryi* occurs also in Sumatra and the Natuna Is.

[Two males were observed courting a female in the jungle at Santubong: the female was situated on a leaf about 12 feet from the ground, over her a pair of males fluttered with a peculiar short up-and-down flight as if they were sliding on perpendicular wires; the female slowly opened and closed her wings but otherwise remained immovable for about three minutes when she suddenly dashed off with the males in hot pursuit, and all three were soon lost to sight].

2. *Hestia belia*, Westw.

*H. belia* Westwood. Cab. Orient. Ent. pl. 37. fig. 2

Sub-sp. *H. belia hypermnestra*, Westw.

Sub-sp. *H. belia belina*, Fruhst.

The typical form of *H. belia* occurs in Java and Sumatra, the two sub-species mentioned above in Borneo, another (*H. bel. linteata*, Butl.) in the Malay Peninsula, and a fourth in Java. Distant (Rhop. Malay p. 406) records *H. linteata*, Butl. from Banjarmassin, but this is probably the same as sub-sp. *belina*.

Genus *Nectaria*.3. *Nectaria leuconoë*, Erichs.

*Idea leuconoë*, Erichs. Nova Acta. Ac. Nat. Cur. xvi p. 283 (1834).

Sub-sp. *N. leuconoë nigriana*, Grose-Smith.

North Borneo, Taganac Island.

Sub-sp. *N. leuconoë chersonesia*, Fruhst.

South Sarawak, Malay Peninsula, Singapore and adjacent islands, Billiton.

Sub-sp. *N. leuconoë natunensis*, Swell.

I am not at all certain that the varieties and local races of this species are all worthy of sub-specific rank; *Natunensis* in especial seems to possess no well-marked characteristics. In the Sarawak Museum collection there is a male from Buntal, Sarawak, which might be either *nigriana* or *natunensis* and a female from Trusan, N. Sarawak is intermediate in character between *nigriana* or *chersonesia*. Other sub-species occur in Japan, Philippines Palawan, Talaut, Sangir, Java, Engano.

The species frequents the sea-shore.

Genus *Ideopsis*.4. *Ideopsis daos*, Boisd.

*Idea daos*, Boisduval, Spec. Gén. Lep. I, pl. 24. f. 3 (1836).

Borneo and Lingga Archipelago, with sub-species n, Singapore, Penang, Malay Peninsula, Sumatra, Nias, China, Hongkong.



The Natuna island forms appears to belong to another sub-sp. *I. daos perakana*, Fruhst.

[This common and distasteful species is mimicked by a Chalcosiine moth, *Isbarta pieridoides*, and by the female of *Papilio delesserti*].

Genus *Danais*.

Sub-genus *Radena*.

5. *Danais* (*Radena*) *vulgaris*, Butl.

*Danais vulgaris*, Butler., Entom. Month. Mag. xi, p. 164, (1874).

A common and widespread species, occurring in Singapore, Malay Peninsula, Sumatra, Java, Nias and Natuna Islands.

[This species and *Parantica eryx*, Fab., are mimicked in Borneo by the following butterflies :—*Elymnias lais*, ♂, *Euripus halitherses*, ♂ and *Papilio megarus*].

6 *Danais* (*Radena*) *juventa*, Cram.

*Papilio juventa*, Cramer. Pap. Ex. ii, pl. 188 B. (1779).

A common sea-side species, ranging nearly all over the Malay Archipelago.

Sub-genus *Tirumala*.

7. *Danais* (*Tirumala*) *septrionis*, Butl.

*Danais septrionis*. Butler, Ent. Month. Mag. vol. xi, p. 163, (1874).

Occurs also in India, Ceylon, Burma, Siam, Malay Peninsula, Java, Sumatra, Formosa.

Mimicked in Borneo by :—*Papilio macareus macaristus*.

8. *Danais* (*Tirumala*) *microsticta*, Butl.

*Danais microsticta*. Butler, Ent. Mouth. Mag. vol. xi, p. 163. (1874).

Occurs also in Java and Nias.

Fruhstorfer considers this to be merely a sub-species of *septrionis*.

Sub-genus *Limnas*.9. *Danaïs* (*Limnas*) *chrysippus*, L.

*Papilio chrysippus*, Linnæus. Mus. Ulr. p. 263, (1764).

[This very widely distributed insect is common in N. Borneo but of extreme variety in the more southern parts of the island; it is interesting to note that the females of *Hypolimnas misippus* which mimic it very closely are hardly ever met with except in N. Borneo though the males are somewhat less rare in the southern part of the island; in other words, the local distribution of the mimics closely follows that of the model].

Sub-genus *Salatura*.10. *Danaïs* (*Salatura*) *plexippus* L.

*Papilio plexippus*, Linnæus Mus. Ulr. p. 262, (1764).

Sub-sp. *D. plexippus intensa*, Moore.

The typical form ranges from Japan through India to the Nicobars, it occurs also in the Philippines. The sub-species *intensa* occurs only in Borneo, Nias and Java; another sub-species is found in Malacca, Singapore and Sumatra.

11. *Danaïs* (*Salatura*) *melanippus*, Cr.

*Papilio melanippus*, Cramer, Pap. Exot. ii. pl. 127 fig. A. B. 1779.

Sub-sp. *D. melanippus hegesippus*.

Fruhstorfer (l. c. 1899 p. 74) has united these two old species. The typical form of *melanippus* he restricts to Java, Nepal, Assam and Penang? (vide Marshall and de Nicéville). The form *hegesippus* occurs in Sumatra, Singapore, Natunas, Malay Peninsula, Penang, Billiton, Mergui Is., and is now recorded for the first time from Borneo. Another sub-species occurs in Burma, Orissa and Bengal.

12. *Danaïs* (*Salatura*) *lotis*, Cr.

*Papilio lotis*, Cramer, Pap. Exot. p. 111, pl. 230, Fig. D. E. (1777).

The typical form is confined to Borneo, sub-species occurring in the Philippines and Celebes.

Sub-genus *Bahora*.

13. *Danaïa* (*Bahora*) *aspasia*, Fab.

*Papilio aspasia*, Fabricius, Mant. Ins. ii, p. 15, n. 145 (1787).

Ranges from Burma, to the Philippines and Banca.

[Mimicked by the female of *Nepheronia lutescens*.]

*D. (Bahora) cleona*, Cram., is a Moluccan species and has been wrongly recorded from Borneo. I cannot agree with Fruhstorfer in regarding *D. aspasia* as merely a sub-species of *D. cleona*.

Sub-genus *Parantica*.

14. *Danaïa* (*Parantica*) *eryx*, Fab.

*Papilio eryx*, Fab. Ent. Syst. Suppl. p. 423. (1789).

Occurs in Borneo with a sub-species in Nias.

Sub-genus *Caduga*.

15. *Danaia* (*Caduga*) *crowleyi*, Jenner Weir.

*Caduga crowleyi*, Jenner Weir. Entomologist, 1894, p. 109.

The species is confined to Borneo, occurring on Mts. Kina Balu and Penrissen. Fruhstorfer considers this to be merely a sub-species of *D. (Caduga) melaneus*, Cr., but this must be quite wrong for *D. melaneus* is a typical *Caduga*, with both patches of scent-scales on the hind-wing traversed by a lengthened swelling of the vein (cf. Moore, Lepidoptera Indica, p. 60). Whilst in *D. crowleyi* only the patch on the sub-median vein is so traversed, the patch on the internal vein is almost obsolete, whilst that on the lower median vein is large and spatulate as in *Parantica*; in short *D. crowleyi* is not a true *Caduga* at all.

16. *Danaïa* (*Caduga*) *luzonensis*, Feld.

*Danaïa luzonensis*, Felder, Wien. Ent. Mon. iv. p. 398. n. 17 (1860).



Sub-sp. *D. luzonensis pramacaristus*, Fruhst.

From N. Borneo: (Mt. Kina Balu) and Mt. Penrissen. The Bornean form has been confused with the Javan *Caduga larissa*, Feld., another sub-sp. of *luzonensis*; other sub-species occur in the Lesser Sunda Is., Sumatra and the Malay Peninsula; the typical form is confined to the Philippines and Palawan.

[The species was common on Mt. Penrissen and often flew in company with *Caduga crowleyi*; *Elymnias lais* ♂ mimicked both species.]

Genus EUPLÆA.

Sub-genus *Menama*.

17. *Euplæa (Menama) lorzæ*, Moore.

*Menama lorzæ*, Moore, P. Z. S. 1883, p. 265, pl. 31, fig 5.

Occurs in North Borneo only.

Sub-genus *Tronga*.

18. *Euplæa (Tronga) crameri*, Lucas.

*Euplæa crameri*, Lucas, Rev. Zool. 1853, p. 318.

*Tronga brookei*, Moore, P. Z. S. 1883, p. 268, n. 8.

*Tronga labuanu*, Moore, l.c. p. 268, n. 9.

*Tronga daatensis*, Moore l.c. p. 268, n. 10.

I agree with de Nicéville & Fruhstorfer in uniting the Labuan and Daat Is. forms with the mainland species *crameri*, and with de Nicéville in sinking *brookei* as another synonym. *E. crameri typica* is confined to the Philippines, Borneo and the Natunas, with doubtful sub-species ranging from India to all the Sunda Islands.

[This and the following species are mimicked by:—*Hypolimnus anomalus* ♀, *Elymnias pellucida*, *Papilio paradoxus telesicles* ♀ ab. *russus* and ab. *leucothoides*, *Papilio leucothoe ramaceus* and the Chalcosiine moth *Isbarta maculurina*.]

19. *Euplæa (Tronga) bremeri*, Feld.

*Euplæa bremeri*, Felder, Wien. Ent. Monat. iv, p. 398, n. 16 (1860).

*Tronga pryeri*, Moore, P.Z.S. 1883, p. 269.

The differences between specimens in the Sarawak Museum of *E. bremeri* (*E. Marsdeni*, syn.) from Singapore and *E. pryeri* from British North Borneo are so small and so inconstant that I have no hesitation in writing the two species under one name. Fruhstorfer divides the *Trongas* into two groups:—

1. Hindwing with a prominent row of submarginal dots.
2. Hindwing with a double series of very large clear white spots.

In the former group he places *E. bremeri*, as a sub-species of *E. Crameri*, in the latter he places *E. pryeri*, yet all the specimens of *E. bremeri* that I have seen should be placed in group 2. In any case the species is rather a doubtful one. It ranges from India through the greater part of the Indo-Malayan region.

Sub-genus *Adigama*.

20. *Euplaea* (*Adigama*) *scudderi*, Butler.

*Crastia scudderi*, Butl. Journ. Linn. Soc. Zool. xiv, p. 297 (1878). Confined to Borneo.

[Mimicked by the *Chalcosiine* moth, *Amesia hyaku*].

Sub-genus *Penoa*.

21. *Euplaea* (*Penoa*) *uniformis*, Moore.

*Penoa uniformis*, Moore, Lepid. Ind. vol. i, p. 99 (1890).

Confined to Borneo; possibly only a sub-species of *E. ulcathor*. God. or *E. ménétrièrei*, Feld.

[Mimicked by *Elymnias lutescens* and *Mimeuplaea tristis*].

22. *Euplaea* (*Penoa*) *zonata*, Druce.

*Euplaea zonata*, Druce, P.Z.S. 1873, p. 338. Confined to Borneo.

[Mimicked by *Papilio slateri hewitsoni*].

23. *Euplaea* (*Penoa*) *masina*, Fruhst.

*Euplaea* (*Penoa*) *masina*, Fruhstorfer, Berl. Ent. Zeit. XLII, 1897, p. 16. Confined to Borneo.

Sub-genus *Trepsichrois*.24. *Euplœa* (*Trepsichrois*) *claudius* Fab.*Papilio claudia*, Fabricius, Gen. Ins. p. 263, (1777).Sub-sp. *E. claudius mulciber*, Cr.

The typical form occurs in India, Burma, Siam, Malay Peninsula, Sumatra and some of the lesser Sunda Island; the sub-species *mulciber* is confined to Borneo, other sub-species occur in Southern India, Java, Nias, Mentawai Island and the Philippines.

[The pupa is a most brilliant object, of a bright burnished gold reflecting like a mirror, with spots and dashes of reddish brown and orange. It is nevertheless not at all conspicuous, since it is always suspended from the underside of a leaf and gives the effect of a hole in the leaf through which the sun is shining. I shall never forget my astonishment, when, on one occasion I attempted to thrust my finger through such a hole and encountered instead the resistance of a large and apparently brilliant pupa.

The *imago* is widely mimicked, the following is a list of the mimics:—

Males. *Euripus halitherses* ♀ *forma cinnamomeus*, *Hypolimnas anomala* ♀, *Elymnias borneensis*, *Papilio paradoxus telesicles* ♂, *Pompeon subcyanea*, *Callamesia striata* ♂.

Females. *Elymnias lais* ♀, *Papilio paradoxus telesicles* ♀, *Callamesia striata* ♀].

Sub-genus *Calliplœa*.25. *Euplœa* (*Calliplœa*) *adyte*, Boisd.*Euplœa adyte*, Boisduval, Bull. Ent. Soc. Fr. 1859, p. 156.Sub-sp. *E. adyte aristotelis*, Moore.

No less than sixteen sub-species of *E. adyte* are distinguished by Fruhstorfer, ranging all through the Malay Archipelago to Melanesia: *aristotelis* is confined to Borneo—*natunensis* occurs in the Natuna Is.



Sub-genus *Macroplæa*.26. *Euphea* (*Macroplæa*) *corus*, Fab.*Papilio corus*, Fab. Ent. Syst. iii, p. 41, (1793).Sub-sp. *E. corus butleri*, Moore, (*Syn. E. Godmani* Moore).

This sub-species is confined to Borneo; the typical form occurs in Ceylon and other sub-species range from Burma through the Malay Peninsula and Sunda Islands to Palawan and Celebes.

Sub-genus *Danisept*.27. *Euphea* (*Danisept*) *diocletianus*, Fab.*Papilio diocletianus*. Fabricius, Ent. Syst. III. 1, p. 40, n. 118 (1793).*Papilio Radamanthus*. Fabricius, Ent. Syst. III. 1, p. 42, n. 127 (1793).Sub-sp. *E. diocletianus lowei*, Butler.

The typical form according to de Nicéville occurs in India, Burmah, Indo-China, Malay Peninsula, Sumatra, Billiton, Banka, Natuna Is. The sub-sp. *lowei* is confined to Borneo. *E. schreieri*, Butler, has been wrongly recorded from Borneo, it is apparently confined to the island of Nias.

[As I pointed out elsewhere \* the females of *E. diocletianus lowei* are extremely rare whilst the males are common; in Singapore both males and females of *E. (Dan.) diocletianus*, Fab., are common. It is at least curious that the female *lowei* should differ noticeably from female *E. diocletianus* whilst the males of both form are almost identical.

The species is mimicked by:—Males:—*Euripus halitherses* ♀ forma *pyciffæra*, *Papilio caunus mendax* ♂,

Females:—*Euripus halitherses* ♀ forma *euplexoides*, *Papilio caunus mendax* ♀ *Mimeurplæa* and also *hadamanthus*].

From Borneo; it occurs in Nias.

---

\*Journ. As. Soc. Straits Br., No. 35, p. 31.

Sub-genus *Salpinx*.28. *Euplœa* (*Salpinx*) *leucostictos*, Gmel.

*Papilio leucostictos* Gmelin (Pap. L.) Syst. Nat. I. 5, p. 2289, n. 889 (1788-1791).

Sub-sp. *E. leucostictos syra*. Fruhst.

Sub-sp. *E. leucostictos kadu*. Esch.

The typical form occurs in Java. I have taken the form *syra* on Mt. Matang. *kadu* occurs in N. Borneo, Palawan and Philippines and is another proof of the Philippine element in the N. Bornean fauna.

Sub-genus *Isamia*.29. *Euplœa* (*Isamia*) *ægyptus*, Butl.

*Euplœa ægyptus*, Butler, P. Z. S. 1866, p. 277.

Occurs in Borneo, with sub-species in Billiton and Philippines including Palawan.

[The species is synposematic with *E. Crameri* and is mimicked by the same species of butterflies and moths].

30. *Euplœa* (*Isamia*) *lowei*, Moore.

*Isamia lowei*, Moore, P. Z. S. 1883, p. 316.

Confined to Borneo, possibly a sub-species of *ægyptus*.

31. *Euplœa* (*Isamia*) *rafflesi*, Moore.

*Isamia rafflesi*, Moore, P. Z. S. 1883, p. 314.

Sub-sp. *E. rafflesi sophia*, Moore.

Borneo and Sumatra, other sub-species occur in Singapore, Nias, and Mentawai, the typical form occurs in Java—De Nicéville (J. A. S. B. vol. lxx, 1901), considers all the above species to be synonymous with *Isamia chloë*, Guér. from Malay Peninsula.

Sub-genus *Stictoplœa*.32. *Euplœa* (*Stictoplœa*) *dufresnei*, Godt.

*Danaïs dufresnei*, Godt. Euc. Méth. ix. Suppl. p. 815 (1823).

Sub-sp. *E. dufresnei tyrianthina*, Moore.

The sub-species occurs only in N. Borneo, where there is a distinct Philippine element. The typical form occurs in the Philippines, other sub-species in Sumatra, Java, Lombok, Sumba, Sumbawa, Alor, Palawan.

Messrs. Pryer and Carter in their list recorded *E. binotata*, Butl., from N. Borneo, but this is certainly erroneous as the species is otherwise only known from N. India. They also record *E. (Stictoplecta) suah*, n. sp. which, however, is a mere *nomen nudum*, no description of the insect being given.

Sub-fam. SATYRINÆ.

Genus *Mycalopsis*.

Sub-genus *Satoa*.

33. *Mycalopsis (Satoa) maineana*, Hew.

*Mycalopsis maineana*, Hewitson, Exot. Butt. iii, p. 87

*Myc.* t. 5, f. 27, 28 (1864).

Occurs also in the Malay Peninsula and Sumatra.

Sub-genus *Orsotriana*.

34. *Mycalopsis (Orsotriana) medus*, Fab.

*Papilio medus*, Fabricius, Syst. Ent. p. 488, (1775).

Ranges over the Indian regions, Malay Peninsula, Sunda Islands, Celebes, Timor, Philippines and Hainan. The dry season form (*runeka*, Moore) does not occur in Borneo.

Sub-genus *Calysime*

35. *Mycalopsis (Calysime) perseus*, Fab.

*Papilio perseus*, Fabricius, Syst. Ent. p. 488 (1775) ♀

Various authors record this species from Borneo, but I have not yet met with it. It occurs in the Indian region, the Malay Peninsula, Sunda Is., Philippines, Hainan and Formosa. The dry-season form does not occur in Borneo.



36. *Mycalesis (Calysime) polydecta*, Cram.  
*Papilio polydecta*, Cramer, Pap. Exot. ii, pl. 144 Fig. e. f. ♀  
 (1777).

In the Sarawak Museum collection is a long series of this species, corresponding very well with the figures 1b, 1d, 1e, 1g. of Plate 61, in Moore's *Lepidoptera Indica*.

The species appears to have been previously recorded from India and Ceylon only; its validity is rather doubtful.

Sub-genus *Culipa*.

37. *Mycalesis (Culipa) mnasicles*, Hew.  
*Mycalesis mnasicles*, Hewitson, Exotic Butt. iii, *Myc.* pl. 5,  
 figs. 32, 33 ♂ (1864).

The species is also recorded from Upper Burma, Tenasserim, Malay Peninsula and Sumatra.

Sub-genus *Martanda*.

38. *Mycalesis (Martanda) janardana*, Moore.  
*Mycalesis janardana*, Moore, Cat. Lep. Mus., E. I. C. i,  
 p. 234, (1857).

Previously unrecorded from Borneo, Dr. R. Hanitsch collected two specimens at Kiou, Kina Balu. Other localities: Malay Peninsula, Singapore (Davison), Java, Sumatra, Celebes (Hose).

Sub-genus *Mydosama*.

39. *Mycalesis (Mydosama) fuscum*, Feld  
*Dasyomma fuscum*, Felder, Wien. Monats. iv, p. 401,  
 (1860).  
 Malay Peninsula, Singapore, Sumatra, Borneo.
40. *Mycalesis (Mydosama) anapita*, Moore.  
*Mycalesis anapita*, Moore, Cat. Lep. Mus., E. I. C. i, p. 232,  
 (1857).  
 Malay Peninsula, Sumatra, Borneo.
41. *Mycalesis (Mydosama) pitana*, Staud.  
*Mycalesis pitana*, Staudinger, Iris. vol. ix, p. 230, 1886.

Jour. Straits Branch

Mt. Kina Balu. I have only seen one specimen captured by Dr. R. Hanitsch at Kiou, Kina Balu.

Sub-genus *Nebdara*.

42. *Mycalasis* (*Nebdara*) *amæna*, Druce.  
*Mycalasis amæna*, Druce. P. Z. S. 1873, p. 339, pl. 32,  
 f. 1. Confined to Borneo.
43. *Mycalasis* (*Nebdara*) *kina*, Staud.  
*Mycalasis kina*, Staudinger. Iris. vol. v, p. 451 (1892).  
 Mt. Kina Balu.

Sub-genus *Suralaya*.

44. *Mycalasis* (*Suralaya*) *orseis* Hew.  
*Mycalasis orseis*, Hewitson, Exot. Butt. iii, p. 83, *Myc.* pl.  
 6, figs. 36, 37, ♂ (1864).  
 Occurs also in Tenasserim, Malay Peninsula, Sumatra,  
 and Nias.

Genus *Neorina*.

45. *Neorina lowi*, D. & H.  
*Neorina lowii*, Doubleday and Hewitson, Gen. D. Lep. p.  
 369, pl. 61, f. 4 (1851).  
 Originally described from Sarawak, sub-species occur  
 in the Malay Peninsula and Sumatra, and in Nias.  
 Mr. W. Doherty (J. A. S. Bengal 1889, p. 124) sug-  
 gests that this species is a mimic of *Papilio helenus* and  
 writes "it may possibly be advantageous for a scarce  
 rather weak-flying insect of *Morphid* or *Satyrin* affinities  
 to resemble a common *Papilio* of powerful and irregular  
 flight"; as far as my experience goes *N. lowi* is much  
 more common than *Papilio helenus*, it is in fact one of  
 the common butterflies of W. Sarawak.

Genus *Celites*.

46. *Celites nothis*, Westw.  
*Celites nothis*, Westwood, Doubleday and Hewitson's  
 Gen. D. Lep. p. 367, pl. 66, f. 2 (1851).

Sub-sp. *Calites nothis epiminthia*, Westw.

The typical form occurs in Siam, *epiminthia* in Borneo, Sumatra and the Malay Peninsula and other sub-species in Borneo and Tonkin.

47. *Calites euptychioides*, Feld.

*Calites euptychioides*, Felder, Reise Nov. Lep. iii, p. 499, (1867).

Borneo, with a sub-species in the Malay Peninsula.

Genus *Lethe*.

48. *Lethe mekara*, Moore.

*Debis mekara*, Moore, Cat. Lep. Mus., E. I. C. i, p. 219, (1857).

Occurs in Sikkim, Assam, Khasias, Burma, Malay Peninsula and Borneo. The dry-season brood does not occur in Borneo.

49. *Lethe cerama*, sp. n.

♂ *Upperside*; very similar to dry season forms of *Lethe mekara*, Moore, from Upper Burma, but of a richer brown and the ocelli on the hind-wing, smaller, but less diffuse. *Underside*, almost exactly the same as in *Lethe delila*, Staud. but the ground-colour is paler and the lilac suffusion less bright; the sub-marginal ocellus on the hind-wing is smaller. Expanse 72 mm.

♀ *Upperside*; forewing, rufous brown merging into fuscous at apex and external margin; hindwing, rufous brown with the abdominal margin pale fuscous and the lower two-thirds of the external margin pale ochreous. The six sub-marginal ocelli of the underside are indistinctly seen on the upperside, the first two as ill-defined black discs, the third as a small black pupil with surrounding ring, the fourth and sixth are hardly visible, the fifth as a large black pupil with surrounding ring. A black marginal line. *Underside*; as in the male, but very much paler, the first ocellus on the hind-wing larger. Expanse 68 mm. *Habitat*: Kuching, Sarawak. Types in the Sarawak Museum.

The male might readily be confused with *L. delila*, Staud. but the female is so very different to the female of that species that I have little hesitation in separating this low country form from the mountain species *L. delila*. The genus *Lethe* is in need of careful revision, a work that is, however, only possible to one who has access to the types of the various species.

50. *Lethe delila*, Staud.

*Lethe delila*, Staudinger, Iris. vol. ix, p. 225, pl. v, f. i. (1896)

Mt. Kina Balu.

51. *Lethe europa*, Fab.

*Papilio europa*, Fabricius, Syst. Ent. p. 500, (1775).

Occurs in the Indian region, Malay Peninsula, Siam, Sunda Is., Philippines, Hainan, Formosa, China.

52. *Lethe perimede*, Staud.

*Lethe perimede*, Staudinger, Iris. vol. ix, p. 226, (1896).

Apparently confined to Borneo.

53. *Lethe darena*, Feld.

*Lethe darena*, Felder, Reise Nov. Lep. iii, p. 498, pl. 68, f. 4.5 (1867).

Sub-sp. *Lethe darena borneensis*, Staud.

The typical form occurs in Java; the sub-species occurs on Mt. Kina Balu.

54. *Lethe dora*, Staud.

*Lethe dora*, Staudinger, Iris, vol. ix, p. 226 (1896)

Borneo only.

Genus *Ypthima*.

55. *Ypthima pandocus*, Moore.

*Ypthima pandocus*, Moore, Cat. Lep. Mus. E.I.C. i, p. 235, (1857).

Malay Peninsula and the Sunda Is.



56. *Ypthima fasciata*, Hew.

*Ypthima fasciata*, Hewitson, Trans. Ent. Soc. (3) vol. ii, p. 287, n. 12, (1865.)

Malay Peninsula, Sumatra and Borneo.

57. *Ypthima abnormis*, sp. n.

♀ *Upperside*; fuscous, without ocelli; the hind-wing is clothed with long hairs especially along the internal margin.

*Underside*—pale brown with dark fuscous striæ which on the forewing are segregated in three areas to form indistinct fasciæ—a sub-basal, a discal and a sub-marginal; on the hind-wing the striæ form five indistinct fasciæ,—a basal, a sub-basal, two discal and a marginal. There are no ocelli. The abdominal margin of the hind-wing is rather deeply excised and the outer margin is slightly sinuate. Cilia fuscous. Expanse, 57 mm. *Habitat*, Kuching, Sarawak.

This is a very aberrant species of *Ypthima*; it has been suggested to me that it is merely a seasonal variation but as I shall have occasion to point out later, the Bornean butterflies do not show seasonal variation; all the species of *Mycalesis*, for instance, correspond to the wet-season phase of the same species, from other countries where the distinction between the fine and wet monsoons is better marked than it is here; the dry-season phases of these species are not found in Borneo.—Type in the Sarawak Museum.

Genus *Ragadia*.58. *Ragadia crisia*, Hubn.

*Euptychia crisia*, Hubner, Zutr. Ex. Schmett. f. 675, 676, (1832).

Occurs in the Malay Peninsula, Penang, Singapore.

The commonest species of the genus.

59. *Ragadia annulata*, Grose-Smith.

*Ragadia annulata*, Grose-Smith, A. M. N. H. 1887, p. 435. N. Borneo.

60. *Ragadia melita*, Staud.

*Ragadia melita*, Staudinger, Iris. vol. v., p. 449 (1892).  
N. Borneo and Kina Balu.

Genus *Erites*.

61. *Erites argentina*, Butl.

*Erites argentina*, Butler, Cat. Satyr. B. M. p. 188, pl. 5, f. 8 (1868).

Upper Tenasserim and Borneo.

62. *Erites elegans*, Butl.

*Erites elegans*, Butler, Cat. Satyr. B. M. p. 147, pl. 2, f. 4 (1868).

Confined to Borneo.

63. *Erites thetis*, sp. n.

♂. *Upperside*; semi-transparent cinereous, on the hind-wings the colour and markings of the underside are seen shining through; a yellow ringed, white pupilled, black ocellus occurs between the second and third median nervules with two much smaller but similar ocelli beyond it.

*Underside*; forewing of same colour as on the upper-side, more transparent at base and this area is crossed by numerous irregular striae; two indistinct ochreous bands cross the wing, one is medial the other post-medial; a row of five sub-marginal small ocelli extends from below 5th sub-costal nervule to the first median interspace, they increase in size from above downwards; a marginal pale band. Hind-wing; basal areas and abdominal margin covered with fine fuscous striae; a medial ochreous band crosses the wing; the outer half of the wing is ochreous; a row of four black ocelli with silvery centres extends from below the 2nd sub-costal nervule to the first median interspace, the lowest of the series is large and has a fuscous suffusion bordering its inner half, the other ocelli are minute; a marginal fus-

cous line. Cilia on forewing cinereous, on hind-wing yellowish-white. The hind-wing is dentate and subcaudate. Expanse 48 mm. *Habitat*: Kuching, Sarawak. Type in the Sarawak Museum.

Genus *Melanitis*.

64. *Melanitis ismene*. Cram.

*Papilio ismene*, Cramer, Pap. Exot. i. pl. 26. figs a. b. (1775).

This common species ranges throughout India, Ceylon, Burma, Andamans and Nicobars, Malay Peninsula, Sunda Is., Hainan, Formosa, Philippines, China and Japan.

[Both the dry-season form *ismene* and the wet-season form *leda* occur in Borneo, but irrespective of the season; I have taken both forms on the same day in the middle of the wet-monsoon and in the middle of the fine monsoon, in fact the two forms fly together. The form, markings, and colouration of the imagines of these seasonal varieties are dependent on the degrees of damp or dryness to which the young stages (egg, larva and perhaps pupa, are subjected, hence a spell of wet weather in the fine monsoon—an event by no means unusual—would produce a brood of wet-season forms and conversely a spell of fine weather in the wet-season a brood of dry-season forms. *M. ismene* in its dry-season phase is remarkably leaf-like, and the insect has the habit of settling amongst fallen leaves and leaning, with both wings closed, over to one side, so that its phyllomorphic appearance is very much increased].

65. *Melanitis zitenius*, Herbst.

*Papilio zitenius*, Herbst, Natursyst. Schmett. viii, p. 5, pl. 182, f. 1, 2 (1796).

This species has previously been recorded only from the Indian region and the Malay Peninsula. The Sarawak Museum collection includes one female in the wet-season phase.

## Sub-fam. ELYMNIINÆ.

Genus *Elymnias*.66. *Elymnias nigrescens*, Butl.

*Elymnias nigrescens*, Butler, P. Z. S. 1871, p. 520, pl. 42, f. 1.

The typical form occurs in Borneo, sub-species are found in the Malay Peninsula, Indo-China, Hainan, Sumatra, Billiton, Lombok, Flores, Sumbawa, Sumba.

This is a non-mimetic species in Borneo, and it is quite the commonest species of the sub-family.

67. *Elymnias hecate*, Butl.

*Elymnias hecate*, Butler P. Z. S. 1871, p. 520, pl. 42, f. 2.  
Confined to Borneo.

This species according to Fruhstorfer is merely a mountain form of *E. nigrescens*, however it is by no means confined to mountains, as it occurs at Labuan and Kuching as well as on Mts. Mulu and Kina Balu and I prefer to look upon it as a distinct and good species.

68. *Elymnias panthera*, Fab.

*Papilio panthera*, Fabricius. Mant. Ins. II, p. 39, n. 40, 407 (1787).

*Elymnias lutescens*. Butler, A. M. N. H. 1867, p. 404, pl. 9, f. 10.

Sub-sp. *E. panthera labuana*, Staud.

Labuan, Sandakan and Kuching, Sarawak.

The typical form occurs in Malacca, Sumatra, Singapore and Natuna Is., sub-species in Java, Banguay, Sulu Archipelago, Palawan, Upper Tenasserim, Nicobars, Nias, Bawean, Engano.

69. *Elymnias dara*, Dist.

*Elymnias dara*, Distant. A. M. N. H. 1887, p. 50.

The male has never been described; a description of



a specimen taken in Kuching follows:—♂ Smaller than ♀ and darker. *Upperside*, dark purplish-black, fasciae on both wings as in ♀ but narrower and shorter and with a lilac tinge.

*Underside*, dark chocolate, the spot on the costal margin smaller than in the ♀, fascia on the fore-wing not so extended. Expanse 57 mm.

The species is confined to Borneo. *E. daedalion*, de Nicév. from Burma is possibly a sub-species. Both *dara* and *daedalion* belong to Moore's sub-genus *Melynius*, not to his genus *Elymnias* as erroneously stated in Lep. Ind. vol. II, p. 154, 155.

70. *Elymnias brookei*, sp. n.

♂. The outer margins of both wings are scalloped; the outer margin of the forewing is produced in the first median interspace into a slight lobe; the outer margin of the hind-wing is produced at the third median nervule to form a short tail. No modified scales on upperside of forewing; androconia on upperside of forewing as in *E. nigrescens*, Butl. *Upperside*: black, on the forewing a sub-apical macular fascia, a few indistinct striae on the costa and an indistinct spot at external angle, blue-green; on the hind-wing an indistinct marginal series of blue-green spots. *Underside*: ground-colour fuscous mottled with dark fuscous, paler along costal margin and apex of forewing and at base of hind-wing, on the hind-wing a sub-marginal series of degenerate ocelli, six in number, black with white centres, the fourth and fifth the largest, the sixth very minute. Cilia on forewing fuscous, on hind-wing fuscous and white alternately, *Antennae* reddish ochreous. Expanse 70 mm. Habitat, Kuching, Sarawak (July). ♀ Unknown. Caught in a trap baited with rotten bananas. In colouration the species approaches *E. esaca* Westw. but the shape of the wings is exactly like that in *E. panthera* Fab. Type in the Sarawak Museum. The species is named after His Highness the Rajah of Sarawak, G. C. M. G.

71. *Elymnias lais* Cram.

*Papilio lais*, Cramer, Pap. Exot. ii, pl. 114, f. A.B. (1779).

Occurs in the Malay Peninsula, Sumatra, Billiton, Java and Borneo, with a sub-species in the Indian region.

[For an account of the habits of this mimetic species see P.Z.S. 1902, p. 259.]

72. *Elymnias pellucida*, Fruhst.

*Elymnias pellucida* Fruhst. Ent. Nach. xxi (1895 No. 11 p. 1) ♀

*Elymnias aroa*, Shelford, P.Z.S., 1902, p. 273. ♂ & ♀

The species is most closely related to *kumara*, Moore: it has been found on Mt. Penrisen, Sarawak, and Kina Balu, N. Borneo.

Messrs. Pryer & Cator record *Elymnias annea* n. sp. from Sandakan, but give no description of it whatever! It may possibly be the same as *E. pellucida*, Fruhst.

73. *Elymnias penanga*, Westwood.

*Melantis penanga*, Westwood, Gen. D. Lep. p. 405 ♀ (1851).

Sub-sp. *E. penanga trepsichroides*, nom. nov.

(*Elymnias borneensis*, Grose-Smith, A.M.N.H. 1892, p. 428.)

There has been much confusion over this species—or sub-species as I prefer to call it. In 1869 Dr. Wallace described (Trans. Ent. Soc. London p. 324,) a female *Elymnias* from Borneo as *E. borneensis*. This species belongs to Moore's sub-genus *Mimadelias* and is a Pierine mimic. In 1887 Staudinger figured (Exot. Schmett. pl. 86) what he supposed to be the male of this species, but Fruhstorfer in 1899 (Berl. Ent. Zeitschr. Bd. xlv p. 57) rightly points out that this is the figure of a female, however he then states that the male of *E. borneensis*, Wall. is "ganz blau und gehört mit *Mehida* Hew. und *Sumatrana*, Wall. zusammen in eine andere

Gruppe und zwar in das sub-genus *Bruasa*, Moore." This is quite wrong, for Grose-Smith in 1892 (l. c.) described both sexes of an *Elymnias* of the sub-genus *Bruasa* from Borneo under the name of *Elymnias borneensis*; the male is blue above and is a mimic of the Euploine butterfly *Trepsichrois claudius*, the female resembles the females of other species of the *Bruasa* section and is not a Pierine mimic as is the female of Wallace's species. Grose-Smith's name *borneensis* being then already occupied by Wallace's species, I venture to propose the new name *trepsichroides*. To make "confusion worse confounded" Grose-Smith described as the female of his *E. borneensis*, the female of another species of *Elymnias* of the section *Bruasa*—*E. konga*—the male of which was described by him in 1899. (A.M.N.H. p. 317.) An undoubted female of *E. penanga trepsichroides* (*E. borneensis* Grose-Smith) from N. Borneo is in the Sarawak Museum collection and is now described for the first time:—Very like the female of *E. penanga*, Westwood, but the sub-apical white fascia on the upper-side of the fore-wings narrower and more outwardly oblique, the costa of the fore-wing striated with white. Underside as in the male but less rufous and darker, the sub-costal primrose-coloured spot, larger than in the male. Expanse 65 mm. Hab. N. Borneo.

74. *Elymnias abrisa*, Dist.

*Elymnias abrisa*, Distant A.M.N.H. 1886, p. 531.

Sub-sp. *E. abrisa konga*, Grose-Smith.

As stated above, Grose-Smith's description of the female of his *E. lorneensis* is in reality the description of a female *konga*; it is very like the female of *E. abrisa*, forma typica, but has more white on the upperside of both wings.

The following is a table of the species and sub-species of the section *Bruasa* of this genus.

*Elymnias penanga*, Westw., forma typica. Penang, Malacca and Singapore (syn. *E. mehidu*, Hew.)

" " *sumatrana*, Wall. Sumatra.

" " *trepsicloroides*, nom. nov. Borneo.  
(=*borneensis*, Grose-Smith.)

*Elymnias abrisa*, Dist., forma typica. Malay Peninsula.

" " *konga*, Grose-Smith. Borneo.

75. *Elymnias esaca*, Westw.

*Melanitis esaca*, Westwood. Gen. D. Lep. p. 405 (1851).

Sub-sp. *E. esaca borneensis*, Wall.

N. & S. Borneo.

Fruhstorfer has done something to clear up the confusion surrounding the species of the sub-genus *Agrusia* (Berl. Ent. Zeitschr. Bd. xlv. p. 56, 57. 1899) but I consider the following table to be a more correct statement of our knowledge of the relationship between the different species and sub-species:—

*Elymnias esaca*, Westw., forma typica. Assam.

" " *borneensis*, Wall. Borneo.

" " *godferyi*, Dist. Malay Peninsula, Sumatra.

" " *andersoni*. Moore. Mergui Archipelago.

" " *leontina*, Fruhst. Nias.

" " nov. sub-spec. (fide Fruhstorfer) Batu  
Is. (Mus. Tring.)

" *maheswara*, Fruhst. Java.

" *egialina*, Feld. Philippines.

*E. esaca*, Westw., has been wrongly recorded from Borneo. Bornean male specimens have a red patch at the base of the hind-wing below, which males of *E. esaca* have not, and I have no doubt at all but that the so-called *esaca* (male) of Borneo is nothing but the male of Wallace's species *Elymnias borneensis* (cf. *antea*). Distant has confused the female of *godferyi* with the females of



Wallace's species and so has wrongly recorded this species also from Borneo ; as pointed out by Fruhstorfer *E. esacoides*, de N., described from a male only, is probably the male of *E. godferyi*, Dist. *E. andersoni*, Moore., I regard as a sub-species of *E. esaca*. The female of *E. esaca* has not yet been described, it will prove to be a Pierine mimic.

Sub-fam. AMATHUSIINÆ.

Genus *Zeuxidia*.

Sect. i.

76. *Zeuxidia amethystus*, Butl.

*Zeuxidia amethystus*, Butler, P. Z. S. 1865 p. 485.

The species appears to have been recorded previously from the Malay Peninsula and Sumatra only.

[In common with nearly all the *Amathusiinae*, this species can be taken in traps baited with rotten fruit].

77. *Zeuxidia doubledaii*, Westw.

*Zeuxidia doubledaii*, Westwood, Gen. D. Lep. p. 329 pl. 52, f. 1 (1851).

Previously recorded from the Malay Peninsula and Penang only.

78. *Zeuxidia wallacei*, Feld.

*Zeuxidia wallacei*, Felder. Reise Nov. Lep. p. 461, pl. 62, f. 3.

Confined to Borneo.

Sect. ii. *Zeuxaltis*.

79. *Zeuxidia pryeri*, Butler.

*Zeuxidia (Zeuxaltis) pryeri*, Butler. A. M. N. H. 1897. vol. 19, p. 469.

N. Borneo

Genus. *Amathuridia*.80. *Amathuridia amythaon*, Doubl.

*Amathusia amythaon*, Doubleday, A. M. N. H. 1847, p. 175,  
*A. amythaon ottomana*, Butl.

The typical form occurs in the Indian region, *ottomana*  
 in Borneo and another sub-species in the Malay Penin-  
 sula.

Genus *Amaridia*.81. *Amaridia aureliana*, Honr.

*Amaridia aureliana*, Honr. Berl. Ent. Zeit. 1889, p. 162.

Confined to Borneo.

This may perhaps be only a sub-species of *A. aurelius*  
 Cr., from the Malay Peninsula and Sumatra.

Genus *Amathusia*.

## Sect. i.

82. *Amathusia phidippus*, Joh.

*Papilio phidippus*, Johanssen. Amoen. Acad. vi. p. 402  
 (1764).

Borneo, Java, Sumatra with sub-species in the Malay  
 Peninsula, Burma, Tenasserim, Nias, Mentawai, Celebes,  
 Torres Straits (?)

83. *Amathusia schönbergi*, Honr.

*Amathusia schönbergi*, Honrath. Berl. Ent. Zeitschr. p. 347  
 T. vi. f. 1 (1887).

Sub-sp. *A. schönbergi borneensis*, Fruhst.

Banjermassin. The typical form occurs in Pegu,  
 Tenasserim and the Malay Peninsula.

Sect. ii. *Pseudamathusia*.84. *Amathusia ochreofusca*, Honr.

*Pseudamathusia ochreofusca*, Honr. Iris. 1886, p. 348.

Malay Peninsula, Borneo, Sumatra.

Genus *Thaumantis*.

## Sect. i.

85. *Thaumantis odana*, Godt.*Morpho odana*, Godart, Enc. Meth. ix. p. 445, (1823).

Malay Peninsula, Nias and the Greater Sunda Is.

Sect. ii, *Kringana*.86. *Thaumantis noureddin*, Westw.*Thaumantis noureddin*, Westwood, Gen. D. Lep. p. 337, (1851).

Malay Peninsula and Borneo.

87. *Thaumantis lucipor*, Westw.*Thaumantis lucipor*, Westwood, Gen. D. Lep. p. 337, (1851).

Malay Peninsula and Borneo.

Sect. iii, *Thauria*.88. *Thaumantis aliris*, Westw.*Thaumantis aliris*, Westwood, Trans. Ent. Soc, 1856, p. 176, pl. 17.

Confined to Borneo.

Genus *Discophora*.89. *Discophora necho*, Feld.*Discophora necho*, Felder, Reise Nov., Lep. iñ, p. 462, (1867).Sub-sp. *D. necho cheops*, Feld.The typical form occurs in Java, *cheops* in Borneo. other sub-species in Sumatra, Nias, Palawan and the Philippines.90. *Discophora tullia*, Cr.*Papilio tullia*, Cramer, Pap. Exot. i, pl. 81, figs. A. B., (1775).

Sub-sp. *D. tullia sondaica*, Boisd.

The typical form occurs in Hongkong, *sondaica* in Java, Sumatra and Borneo, other sub-species in India, Tenasserim, Malay Peninsula and the Philippines.

91. *Discophora amethystina*, Stich.

*Discophora amethystina*, Stichel, Berl. Ent. Zeitschr. xvi. S. B. p. 4, (1901).

Borneo. I am not acquainted with this recently described species.

Messrs. Pryer and Cator also record *Discophora celinde*, Stoll. and *Discophora ogina*, Hübn, from Borneo, but without having actually taken specimens of these species, so that the records must be regarded as extremely doubtful.

Genus *Enispe*.

92. *Enispe milvus*, Staud.

*Enispe milvus*, Staudinger, Iris vol. ix, p. 231, pl. v. f. 4, (1896).

Mount Kina Balu.

Marshall and de Niceville in Butterflies of India, vol. i, p. 312, record *Stichophthalma nourmahal*, Westw. from India, Sikkim and Borneo; the latter locality is evidently erroneous and I can find no confirmation of it in any other publications on Oriental butterflies.

Genus *Clerome*.

93. *Clerome phaon*, Erichs.

*Papilio phaon*, Erichson, N. A. Acad. N.C. p. 401, pl. 50, figs 1, 1a (1834).

Philippines and Borneo.

Westwood (Trans. Ent. Soc., London, 1856, p. 186), gives Borneo as one of the localities of this species, the locality has up to the present never been confirmed; there are however several undoubted specimens of this



species in the Sarawak Museum collection from Limbang and Trusan, N. Sarawak. As before mentioned there is an infiltration of Philippine forms to be discerned in the N. Borneo fauna.

94. *Clerome gracilis*, Butl.

*Clerome gracilis*, Butler A.M.N.H., 1867, p. 401, pl. 8, f. 7.

Malacca, Singapore, Borneo, Sumatra.

I cannot agree with Fruhstorfer in regarding this as a sub-species of *phaon*.

95. *Clerome stomphax*, Westw.

*Clerome stomphax*, Westwood, Trans. Ent. Soc., 1856, p. 186, pl. 21, figs. 3, 4.

Borneo.

96. *Clerome besa*, Hew.

*Clerome besa*, Hewitson, Exot. Butt. iii, *Cl.* pl. 1, fig. 1, (1863).

Fruhstorfer considers this to be merely an aberration of the preceding species.

Borneo.

97. *Clerome arcesilaus*, Fab.

*Papilio arcesilaus*, Fabricius, Mant. Ins. ii. p. 28, (1787).

Indian region, Siam, Malay Peninsula, the Greater Sunda Is. and Bali.

98. *Clerome kirata*, de Nicév.

*Clerome kirata*, de Nicéville, Journ. Bombay Nat. Hist. Soc. 1891, p. 344, Pl. F. fig. 3.

Malay Peninsula, Borneo (Kina Balu), Sumatra.

Genus *Tenaris*.

99. *Tenaris occulta*, Grose-Smith.

*Tenaris occulta*, Grose-Smith, A.M.N.H. 1889. p. 316, Borneo.

Genus *Xanthotzenia*.100. *Xanthotzenia busiris*, Westw.

*Xanthotzenia busiris*, Westwood, Trans. Ent. Soc. London, 1856, p. 187.

Tenasserim, Malay Peninsula, the Greater Sunda Is., and Nias.

Genus *Amnosia*,101. *Amnosia baluana*, Fruhst.

*Amnosia baluana*, Fruhstorfer, Ent. Nachr. xx, No. 19, p. i, (1894).

N. and S. Borneo.

Herr Fruhstorfer informs me that in his collection is a female of this species from S. Borneo which differs somewhat from the type female from Kina Balu; I have only seen specimens from Mt. Matang near Kuching and these do not appear to differ in any way from the published description of the Kina Balu form: Herr Fruhstorfer's S. Borneo specimen is evidently from the low-country.

The position of this genus is very doubtful. I follow Fruhstorfer in placing it amongst the *Amathusiinae*; de Nicéville suggested that it should come at the end of the *Satyrinae*, whilst Felder and Schatz-Rüber placed it amongst the *Nymphalinae*, probably its correct position.



## The Sakais of Batang Padang, Perak.

BY G. B. CERRUTI.

The word Sakai is the Malay name for the aborigines who inhabit the forest on the high slopes of the lower half of the main ridge and some subsidiary ridges of mountains of the Malay Peninsula. As the Malays were the first to come into intercourse with these aborigines, the influence of the Malay, as well as the fear of them, is strong upon them. Malay history in Perak reaches with certainty no farther back than the 16th century, and Malays have no written records relating to the Sakais, whom they treated as slaves and less than human beings.

The Sakais themselves have neither written records nor signs to represent language, whatever information regarding their origin is supplied by them rests solely in tradition.

The narrative of events, which is extracted from them with difficulty, very seldom passes beyond the time of a grandfather, and may be regarded as inexact if not incoherent.

We shall, therefore, have to look for knowledge of their origin to the results of a morphological study of the race.

### PHYSICAL CHARACTERS.

The average height of the male Sakai may be taken approximately at 5 ft. 3 in. : and that of the female at 4 ft. 11 in. : These figures are for the present only tentative. The colour various from a light to a chocolate-brown, the eyes are slightly almond shaped, the nose is flat, the forehead straight, the lips full and separate, but not negro like, the teeth regular and well-formed though blackened by sireh, the hair copious, black, somewhat wavy, occasionally crisp, but never woolly. The senses are unusually keen and well developed. In his native jungle he sees better, hears better, and apparently uses his sense of smell better than other races. His touch is delicate and sensitive, as is that of most savage races, and his sense of



taste is his criterion to judge of the good or ill effects of many objects.

The body and limbs are generally speaking well formed. Cripples and deformed children are extremely rare amongst the Sakais, nor are abnormalities of anatomical structure frequent amongst them.

#### MENTAL AND MORAL CHARACTERS.

A desire for what may be called independence, but what in reality is a dislike of restraint is remarkable in this race. Work for a Sakai must be voluntary; the moment that it becomes compulsory it becomes distasteful.

Not less notable is his distrust of strangers. The approach of a white man will often scatter a whole habitation of Sakais; and even the presence of natives of other races, such as Malays, Tamils or Chinese, is a frequent cause of their speedy removal from an accustomed haunt. Once the Sakai confidence is secured, he is like a child, and must be treated as such. All obligations entered into with him must be scrupulously observed, for, like the natural child, he is not prone to deceit or falsehood. He is also possessed of the child's simple idea of morality, as expressed in his words and acts. Early marriage being the custom, the immorality of civilized races, with its literature and influence on social relations, is unknown.

#### DRESS.

Bark beaten finely and elongated until it resembles coarse ramie fibre, is the material from which the primitive clothing is made. Both sexes fasten strings of this bark, about six inches or more wide, around the waist, by tying them in back and in front. A thin fillet of the same stuff, dyed and coloured in a simple pattern, is used to tie the hair, which is generally filled by the women with combs, made of bamboo and ornamented in various styles.

Flowers are universally worn by the women in the hair, around their necks as necklaces, and occasionally in their waist-belts of odorous grass. In both sexes the nasal septum is perforated for the insertion of straight pieces of bamboo, and the ear only by the women for the insertion of some bamboo with

some odorous grass, of shell, and of animal teeth, to serve as ornaments.

#### ORNAMENTS.

The use of necklaces belongs to attire. Ornamentation of the body is effected by painting the skin in different colours, mostly red, yellow and black, by dyes obtained from plants, gutta and lime. Two lines, one drawn from the vertex of the head over tip of nose to chin, and the other from ear to ear, bisecting the first, divide the face into four areas, the painting of two of which on one side must correspond to the painting of the two others of the opposite side.

The chest and body are generally divided also by a vertical line cleaving the trunk in two halves, right and left, upon which similar patterns are painted.

The object of this adornment by painting is not merely decoration, it is what formerly would have been called dedicated to superstitious uses. The painting of the face and body is, in fact, a species of charm and is supposed to act as amulets and talismans are presumed to act, by warding off dangers, driving afar evil spirits, and filling the wearers with unusual courage.

#### RELIGIOUS SENTIMENT.

It is a peculiarity of the Sakai that, like many of his characteristics, his religious belief is extremely simple. The idea of a Creator, of an all-powerful, all-just and all-merciful Ruler, is absent from his scanty mythology. The origin of the world and the life of mankind on the earth present no problems to him. He believes simply in good and evil spirits. The good spirits are to him vague, indefinite beings, who manifest themselves rarely fatally, and about whom, therefore, he knows and believes little. The evil spirits, on the contrary, are feared, because they are considered to dwell in dangerous ravines, in abandoned *kampongs*, in caverns, and in places regarded according to popular Sakai report as uncanny, whence they issue to infect the Sakais with famine and disease. They also are believed to make themselves felt in thunder, in lightning, and most particularly of all in wind. The early morning breeze

which blows on the tropical hills after 2 a.m. is for the Sakai the work of bad spirits. It is the hour when he feels the fall of temperature most, and it is for him a fatal hour. Accordingly, all true Sakais are awake at this time to chat and smoke and wait until the biting blast passes away. Tattooing is little known amongst them; and though they have a certain knowledge of the ways of tigers and snakes, these are neither worshipped nor considered to be directly concerned with evil spirits.

#### SOCIAL RELATIONS.

The most important circumstance of a man's, and of woman's life, marriage, does not loom large in the Sakai's mind. For him it is neither a religious ceremony nor a civil contract, it is merely a mode of sexual union founded upon mutual sympathy. Rites in connection with it have so far not been proved to be practised. There is neither capture, nor purchase, nor selection. The elders, moreover, do not appear to interfere in the choice of their sons and daughters.

It is probable that it is owing to this fact that these aborigines are gradually decreasing in numbers, even though consanguinity in matrimonial relationship is forbidden, about the only prohibition of any kind that the Sakais know, and to which they submit.

Polygamy exists, but it is rare. Divorce also exists, and is common. The marriage tie, being as loose as is described, is unable to consolidate a union; the slightest incompatibility of temper, temporary sterility of the wife, lasting about three durian seasons, or an attack of dangerous disease, is a sufficient cause for a divorce, which is accomplished without resentment or apparent jealousy on either side. Deformed persons which are very rare amongst the Sakais, or those attacked by dangerous disease, must make a vow of celibacy. The women give birth to their children with only old women attendants, but in a place prepared by the husband. The child is not allowed to touch the earth, either from a superstition that the child would be injured by contact with the earth, or that the child would soil the ground, but is laid upon a couch of dry leaves, which cover a rudely made clay embankment. Directly after birth only old women and young children who are not able and



strong enough to enter the jungle to find their daily food are permitted to approach the child. All others are excluded for a certain period, as there is a certain superstition among them that able bodied persons approaching a newly born baby will contract its smell and take it to the jungle with them when out looking for food. The evil spirits, it is said, are always on the look-out for persons with this smell, and will follow them on their return to their huts to the birth place of the child. At the end of that time the child receives what may be called a ceremonial purification of water, and is presented to him at the village.

#### HABITS.

The Sakais are essentially nomadic, and clear only very limited areas in the hill forests for cultivation; of rice culture they know little, for corn or maize and the Sikoi, sweet potatoes, and tapioca, are their principal crops. The most primitive of the Sakais still subsist by the chase, using the Sumpitan, or blow-gun, and poisoned darts to kill wild animals and birds. As is well-known, the darts are poisoned by being dipped in a gummy or glutinous extract of Ipoh which hardens on the tips, and of another and more dangerous poison extracted from the roots of a kind of creeper named by the Sakais *Legop*.

The Sakai dies as he lives, surrounded by powers of nature which he understands not. If a disease be regarded as contagious, a noise is made on rude drums made of big bamboo to drive away the evil spirits. It is remarkable that there are not musical instruments to express grief; but in expression of joy a flute played through the nose, and a kind of mandoline made also of bamboo, are performed upon particularly by women. After death comes burial in a deep grave, the body generally standing erect in the grave about 4 feet deep or in a sitting posture with tobacco, betel-nut, potatoes, fruits and also with his blow-pipe and poisoned darts by his side. The grave is closed by felling some jungle surrounding it and for about a week they bring the usual food, if a female also some flowers, and afterwards abandon the neighbourhood; for a dead person frequently drives the timid Sakais miles away from promising slopes on which they were beginning to grow their necessary food.





## On Some Hymenoptera From the Raffles Museum, Singapore.

BY P. CAMERON.

Dr. Hanitsch having sent me to be named some undetermined Hymenoptera from Singapore, I give a list of them as a small contribution towards the knowledge of the Hymenopterous Fauna of the Island.

*Evania appendigaster* Lin. A cosmopolitan parasite in the egg-cases of Cockroaches.

*Stilbum splendidum*, Fab.

*Macromeris violacea*, Lep.

*Discolia decorata*, Burm.

This species (which = *D. flavopicta* Lm.) is in the collection of the Raffles Museum from the Dindings.

I think it very probable that *D. ergenna*, Com. (Journ. St. Br. Royal Asiat. Soc., 1902, p. 82) is its male.

This species is recorded by Magretti (Ann. ch. Museo Civico di Storia Nat. di Genova (2) xii, 243) from Schwegoo, Burma; but it is not included by Bingham in the Fauna of British India, Hymen. It has been reported from Java and Sumatra.

*Salix flavus*, Fab.

*Sceliphron violaceum*, Fab.

*Irypoxylon petiolatum*, Sm. Found in the Museum Workshop.

*Piagetia ruficollis*, sp. nov.

Black the scape of the antennæ, the greater part of the clypeus, the prothorax, the mesonotum in front of the tegulæ, the tegulæ, the abdominal petiole, except at the base, and the

legs, red; the 4 front coxæ above, the hinder entirely, the basal point of the 4 front, trochanters, of the hinder above, a line on the fore femora behind, the apical two thirds of the hinder above, the greater part of the hinder tibial and the base of the hinder coxæ, black. Wings hyaline, the basal half of the radial cellule, the apex of the 1st cubital cellule, the greater part of the 2nd and 3rd and the discoidal along the recurrent nervure smoky; the nervures and stigma black.

Head and thorax covered with silvery pubescence ♀

Length nearly 10 mm.

Hab. Singapore, June.

Clypeus indistinctly keeled down the centre; there is a semi-circular depression in the middle at the apex, which has a distinct margin and has a slight incision. Base of mandibles broadly yellowish testaceous. The base of the hinder femora is slightly thickened below, the apex of the thickened part ending in an indistinct tooth.

Comes nearest to *P. rufivenis*, Cam. which may be known from it by the antennæ being almost entirely red, and the sides and apex of the median segment are also red.

*Rhyrchium hæmorrhoidale*, Fab.

*Vespa cincta*, Fab.

*Icaria Singaporensis*, sp. nov

Brownish-black, the head below the antennæ, except for a narrow black line down the centre, a mark, twice longer than wide and dilated above, a mark twice longer than broad, transverse above, narrowed and rounded below, on the front the eye incision and the lower inner orbits broadly, the upper orbits narrowly, the line dilated above to the hinder ocelli, the outer orbits entirely below, the inner half of the upper part, the mandibles, except the teeth, a line on the pronotum, the base and the lower half of the propleuræ, the meso and metapleuræ, except for an oblique black line on the former and extending from the lower furrow downwards and with a short line on either side of its top, 2 lines on the mesonotum, the base of the scutellum, the post-scutellum, 2 large lines on the centre of the

metanotum, the sides of the petiole to near the apex, 2 round marks on the centre of the post-petiole, the extreme base of the 2nd segment, its sides to shortly beyond the middle broadly, the mark at the apex diverging towards the middle of the segment, 2 large marks on the basal half of the 3rd, the marks broader than long and rounded on the inner side, the 2nd segment below, except along the sides and apex, the latter with the sides broadly and roundly dilated and the centre transverse, the base of the 3rd segment, the line narrowed and transverse in the middle and lines on the sides of the apical segments, yellow, legs black, all the coxæ, the lower side and the apical half of the femora above, the underside of the tibiæ and their apex above, yellow; the apical joint of the fore tarsi of a more obscure yellow. Wings hyaline, with a violaceous-tinge, the nervures and stigma black.

Length 14 mm. ♀

Hab. Singapore.

Scape of antennæ below yellow, the flagellum beneath and its apex above, rufous. There is an indistinct keel on the lower part between the antennæ. The black on the front is tinged with rufous. Thorax smooth, the scutellum closely, minutely punctured, its apical half furrowed in the centre. The petiole is longish as long as the 2nd and 3rd segments laterally together; the dilated apex is somewhat twice longer than wide. Head broader than the thorax.

Comes near to *I. 4-maculata*, Cam. The present species is more slenderly built and with a more slender petiole in particular being more slender and not dilated in the middle.

*Icaria rufinoda*, sp. nov.

Deep black, densely covered with white pubescence, the apex of the clypeus and the base of the mandibles pallid yellow, the petiole ferruginous; the wings hyaline, the whole of the radial cellule and the greater part of the apex from the 2nd transverse cubital nervure smoky, with a violaceous tinge, the nervures and stigma dark fuscous. ♀

Length 12 mm.

R. A. oc., No. 41, 1904.



Hab. Singapore, June.

Front and vertex alutaceous, the face and clypeus densely covered with a white pile. Thorax alutaceous, covered with a white pile. Metanotal furrow deep, the sides oblique, the bottom with a narrow smooth impression; it is not striated. Scutellum and post-scutellum coarsely alutaceous, almost rugose; the apical slope of the post-scutellum smooth and shining. Abdominal petiole nearly as long as the 2nd segment, the basal third narrowed; the basal half of the dilated part obliquely narrowed towards the base, the 2nd segment bell-shaped, its length slightly greater than its width at the apex, which has a distinct crenulated furrow, the apical segments with a silky pubescence. Legs primrose, the spurs black.

Comes close to *I. lugubris* Sm. which may be known from it by the black abdominal petiole.

*Nomia iridescens*, Sm.

*Crocisa emarginata*, Lep.

*Anthophora zonata*, Lin.

---

### Malay Hymenoptera Addenda and Corrections

In my paper (J. S. B. R. A. No. 39, 1903) I have omitted to state that Mr. Shelford reared *Spinaria curvispina* Cam. from the larva of a species of *Thosea*, a moth of the Family *himacodidæ* and *Dedanima longicornis* Cam. from a species of *Charocampa*.

I take this opportunity of adding the descriptions of two new Malay species of *Bracon*.

*Bracon teius* sp. nov.

Black; the head pallid yellow; the pro. and mesothorax and the sides of the median segment on the basal half, ferruginous; the 4 front legs ferruginous, the middle tarsi fuscous, the hinder legs black, thickly covered with black hair, the calcaria dark testaceous. Wings dark fuscous, violaceous, with an indistinct hyaline, oblique cloud in the 1st cubital cellule. ♀

Jour Straits Branch

Length 16; terebra 20 mm.

Hab. Ternate.

Face rugose, covered with long pale hair. Apex of 1st abdominal segment closely, distinctly longitudinally striated; the plate on 2nd segment clearly longer than its greatest width irregularly striated in the centre, its keel reaching to the base of the apical third of the segment; from its outer side a keel runs obliquely to the apex; the part between the keels bears curved, oblique, clearly separated striæ; the securiform articulation and the furrow on the next segment striated; in the centre the striæ are continued on to the centre of the segment. The 2nd and 3rd abscissæ of the radius together are about equal in length to the 1st. The abdomen is narrow not dilated in the middle and is clearly longer than the head and thorax united.

*Bracon spilogaster*, sp. nov.

Black, the head pallid yellow, the thorax and 4 front legs ferruginous; the wings fuscous, the stigma and nervures black. ♀

Length 9 mm.; terebra 4 mm.

Head smooth and shining; the face and vertex covered with fuscous pubescence. Thorax smooth and shining; the metanotum has 2 blackish marks on the apex. Abdomen as long as the head and thorax united; black, the basal 4 ventral segments pale, with 2 large black marks in the centre; broad in the middle, narrowed at the base and apex; smooth and shining; the securiform articulation stoutly striated; the keel on the 2nd segment is longer than broad, is broad at the base, becoming gradually narrowed towards the apex, which is prolonged into a short keel with a depression on either side, but not reaching to the apex of the segment; the furrows on the 3rd and 4th segments are narrow, curved and smooth.

The tibiæ and tarsi are thickly covered with a pubescence and, more sparsely, with pale hair; the hinder calcaria fuscous; the 2nd cubital cellule in front is about one-third longer than the 3rd. Mandibles pale yellow, the teeth black.

*P. Cameron.*

## ERRATA.

In my paper in the "Journal" for 1902, No. 37 occur the following printer's errors.

- Page 30, 13th line from bottom for "covered" read "curved"  
 " 31 & 32 for "Megiselens" read "Megischus."  
 " 33, 10th line from top for "sharpened" read "shagreened"  
 " 34, 1st " " for "smoothy" read "smoky."  
 " 37, 17th " " for "Brule" read "Brulle."  
 " 39, 5th " " for "expressed" read "depressed"  
 " 39, 16th " " for " " read " "  
 " 44, 7th " " for *acvenitini* read "*acoenitini*."  
 " 44, 10th " " for *acvenites* read *acoenites*.  
 " 47, 11th " " for Fah. read Fab.  
 " 50, 3rd line from bottom add after "smooth" *peronatum*.  
 " 50, add after "reticulated" in last line *fuscicorne*.  
 " 51, above *anisobas cincticornis* add *Ichneumonini*.  
 " 52 " *Bodargus* add *Joppini*.  
 " 53 " *Diapetus* add *Cryptini*.  
 " 62 2nd line from bottom for "slope" read "shape"  
 " 71 14th " top *Joppini* should be placed above *Zono-*  
*joppa*.  
 " 73, 5th " bottom for "are" read "areæ."  
 " 81, top line for "metapleurg" read "metapleuræ."  
 " 91, 9th line from top for "sharpened" read "shagreened"  
 " 114, 12th " bottom for "tubæ" read "tibiæ"  
 " 125, 2nd " " for "tech" read "teeth"  
 " 138, 3rd " " for "covered" read "curved"  
*P. Cumeron.*

## Correction to Journal No. 39.

Page 54 after line 18 insert.

"Follow the principal noun with which they are connected; and the object."

## Short Notes.

---

### On the Flowering of *Barringtonia racemosa*.

The *Barringtonias* are trees of moderate size, belonging to the order *Myrtaceæ* and usually to be met with along tidal rivers, or more rarely in the hill woods. The flowers are produced in long hanging racemes, or in some species in short erect spikes. In *B. racemosa* the pendulous spikes are about  $2\frac{1}{2}$  feet long and bear about 30 flowers. They are sessile with a short  $\frac{1}{2}$  inch ovary with 2 or 3 rounded green sepals and four lanceolate white petals, an inch long. The stamens are innumerable, with slender filaments an inch long and minute yellow anthers, the style is nearly as long slender with a minute capitate stigma and all deep crimson.

The peculiarity of its flowering consists in the fact that it is nocturnal. The flowers open about 4 or 5 to 16 on each spike at a time, the buds commence to split about mid-day, and remain partially open till nightfall, fully expanding at about half-past seven or eight. At that time the petals are spread out widely, and the stamens radiate in all directions, so that the flowers have a brush-like appearance.

Before daylight the petals and stamens have fallen in a mass, leaving only the calyx and the stiffly projecting style.

They exhale a rather coarse scent somewhat resembling that of meadow-sweet, and from a tree with about 20 flowers open at once I could perceive the scent distinctly at 25 paces off. The honey, abundant at night, is contained in a nectary formed by the connate bases of the stamens. The flowers are visited by moths, I caught a common grey Noctuid, and a rather curious looking light red brown *noctua* with plicate wings.

Moths were not very abundant at the flamers, when I observed them, but perhaps this was due to the strong moonlight.



The smaller brown moth plunged into the flower among the stamens so as to reach the honey.

I have little doubt that a large tree of the genus *Careya* (apparently undescribed) in which the flowers were of similar shape but borne in an erect short spike, is fertilized in the same way, as though in full flower no open flowers were procurable during the day time, but the whorls of stamens were found covering the ground in the mornings. In this lofty tree in the Botanic gardens jungle the stamens were white but the base of the filaments crimson.

The Myrtaceæ as a rule seem to be day flowering plants. The *Eugenias*, our biggest genus, have usually white flowers often produced in large corymbs. *Eugenia lineata* and similar species are haunted, when in flower, by abundance of bees. *Apis dorsata* and *A. florea*, *Trigona collina* and other species and the pollen-eating flies (*Syrphidæ*) and also by many butterflies.

*E. Ridleyi* peculiar from the flowers being light apple green in colour is visited by flies (*Muscidæ*).

*Rhodamnia trivervia* with small white sweet scented flowers produced in great abundance and lasting but a day each, is visited by bees, *Apis*, and *Trigona*, and by the *Syrphidæ*.

H. N. Ridley.

#### Fertilization of *Webera Stellulata*.

*Webera Stellulata* Hook. fil. is a small shrub 2 or 3 feet tall belonging to the order *Rubiaceæ*. It has smooth dark green shining leaves elliptic cuspidate, and a short dense corymb of light green flowers. The buds are peculiar in shape, being fusiform and narrowed towards the tip, the joints of the petals instead of being pressed together at the top into a point as in the other species of *Webera* are turned out to one side bent at an angle pointing from left to right. They are green and covered with white hairs and at the base they are connate into a short tube, in the mouth of which are long white hairs. The stamens five in number have short green filaments and long linear anthers, which split and shed their pollen before the

Jour. Straits Branch

flower opens. The style is long and cylindrical and covered entirely with white hairs. When the fully developed bud is touched on the tip, the petals suddenly spring open and lie quite flat in the form of a star. At the same time the pollen lying loose in the bud is thrown upon the other flowers already open. The mechanism by which this sudden expansion of the flower takes place seems to be very simple. The upper part of the petals are twisted in bud, and on the side opposite to the direction in which the bent tip points a portion of the edge is incurved so as to be tightly held by the next petal to it. A light pressure, as of an insect, on the horizontal tips of the petals by bending them down causes the petals to separate and fly back suddenly, jerking the pollen out over the other flowers, or possibly on the insect visitor. On the top of the ovary is a brown sticky ring which may perhaps secrete honey but I cannot detect any in the tube. The flowers possessed a faint scent, and may be fertilised by insects, but considering the inconspicuousness of the green flowers, as compared with the sweet-scented white blossoms of the other *Weberas* and the fact that it is quite easy for the pollen of one flower to be thrown by the mere opening of the flower on to the stigma of the adjacent one, it is more than probable that the plant itself fertilises one flower by the pollen of another.

*Webera stellulata* inhabits forests, usually in rather dry spots. I have found it in Singapore at Woodlands flowering in June, and Bukit Mandai, also in Johore, at Panchur; Selangor, Kuala Lumpor; Negri Sembilan, Gunong Angsi.

The Malays call it Kahwa hutan and Kuruseh putih and Pokoh Subiroh. It flowers from December to June.

H. N. Ridley.

### Human Images among the Orang Mantong.

I have long suspected the existence of *Berhala*, or human images, among the "wild tribes" of the Rhio archipelago, but never actually met with any until the past summer (1903).

When at Pulo Sanglar or Lake Durian, Rhio archipelago, in July. I found two wooden images representing women, in a cave near the sea shore, not far from Kampong Telok Lanun.

Each image is about  $3\frac{1}{2}$  feet high. One of harder wood was much more carefully carved than the other. It had 3 wooden horns about 6 inches long projecting upwards from the head. These horns were serrated on one edge. This figure also had straight rudely carved arms of soft wood, much decayed.

The teeth were represented by pieces of broken shell. A blackish line extended diagonally across the chest, meeting a horizontal line extending across just above the position of the nipple. A blackish spot was over the position of the heart.

The other figure was very rudely carved of soft white wood and was without arms.

The figures were lying face downward on the floor of the cave and had evidently not been disturbed for months, as roots were growing over them and the wood was beginning to decay.

Pulo Sanglar is inhabited by Orang Mantong, and latterly many Chinese have settled there cultivating gambier.

The Batin of Telok Lamun called himself a *Malay*, but he was more than half Orang Laut.

No true Malays live on Sanglar, but they inhabit the neighbouring small islands. These Malays call all the Sanglar people *Tambus*, except of course the Chinamen, and say there is not a Mussulman. All the Sanglar people eat pig. They are certainly not true *Tambus*. They were very shy, and I had a lot of trouble inducing them to be photographed.

No information in regard to the use of the images could be obtained. Every one denied the existence of such things, not knowing I had already found them.

The images cannot be regarded as true *berhala* or idols. Most probably they are a sort of "*Sakkat buang*" for use in



sickness. Among the Orang Laut when a man is ill, a wooden figure of a bird, snake, fish or other animal is made, and the *pawang* or *bomo* exorcises the hantu or devil in the sick man and drives it into the figure, which is then carried out to sea and thrown overboard. Last year we picked up a wooden bird floating in Durian Strait.

Very likely the human figures were used in the same way, being carried out into the jungle instead of out to sea. Like the Rumah hantu to be seen in the woods near Malay Kampongs. These images resemble the adu adu of Pulo Nias.

*Dr. Abbott.*

### The Orang Laut of Singapore

In Journal 33, p. 247, Mr. Skeat and I published some notes on the Orang Laut of Singapore, a race very nearly extinct, and of which very little is known, I have since come across an account of them in Finlayson's Mission to Siam and Cochin China, in 1821. The author somewhat naturally mistook them for Malays and thus describes them. "The condition of the lower class of Malays in these parts is wretched beyond what we should conceive to be the lot of humanity in an intertropical climate, almost the whole of their life is spent upon the water in a wretched little canoe in which they can scarce stretch themselves for repose. A man and his wife and one or two children are usually to be found in these miserable sampans; for subsistence they depend on their success in fishing. Their tackling is so rude and scanty that they are often reduced to the most urgent want, when they have made a meal they lay basking in the sun or repose under the dense shade of the mangrove till hunger again calls them into action. They have scarce a rag of cloth to secure them from the scorching noon-day sun or shelter them from the damp and noisome dews and exhalations of night. The women are not less dexterous than the men in managing their boats. Their only furniture consists of one or two cooking pots, an earthen jar and a mat made of the leaves of the Pandanus which serves to protect them from the rain. In the numerous bays inlets and creeks that surround Singapore an inconceivable number of families live in



this wretched manner who have never possessed a house nor any sort of abode on the land. They are constantly roving about from place to place in pursuit of fish. What they have succeeded in taking more than is required for their immediate use, they dispose of to the fixed inhabitants, taking rice, sago, betel and cloth in return. This description of Malays goes by the appellation of Orang Laut or men who live on the sea.

A number of the people called Orang Laut were brought to us for inspection. They were superior in condition, in appearance more civilized than many whom we had seen in the bays and creeks remote from the haunts of men. A portrait was taken of one of them illustrative of the physiognomy and general appearance of the Malay race, six of these men were more minutely examined. Their average height was five feet three inches, average weight nine stone eight pounds, average circumference of the chest two feet ten inches, circumference of the clenched fist about eleven inches, average of facial angle  $66\frac{1}{2}$ , average temperature under the tongue  $100\cdot02$ ."

*H. N. Ridley.*

